




The Georgia Agricultural Experiment Stations
College of Agricultural and Environmental Sciences
The University of Georgia

Annual Publication 104-3
January 2012

GEORGIA

2011 Peanut, Cotton and Tobacco Performance Tests

J. LaDon Day, Anton E. Coy, Stevan S. LaHue,
Larry G. Thompson and John D. Gassett
Editors



Department of Crop and Soil Sciences
Griffin Campus

Conversion Table

U.S. <i>Abbr.</i>	<i>Unit</i>	<i>Approximate Metric Equivalent</i>
Length		
mi	mile	1.609 kilometers
yd	yard	0.9144 meters
ft or '	foot	30.48 centimeters
in or "	inch	2.54 centimeters
Area		
sq mi or mi ²	square mile	2.59 square kilometers
acre	acre	0.405 hectares or 4047 square meters
sq ft or ft ²	square foot	0.093 square meters
Volume/Capacity		
gal	gallon	3.785 liters
qt	quart	0.946 liters
pt	pint	0.473 liters
fl oz	fluid ounce	29.573 milliliters or 28.416 cubic centimeters
bu	bushel	35.238 liters
cu ft or ft ³	cubic foot	0.028 cubic meters
Mass/Weight		
ton	ton	0.907 metric ton
lb	pound	0.453 kilogram
oz	ounce	28.349 grams
Metric <i>Abbr.</i>	<i>Unit</i>	<i>Approximate U.S. Equivalent</i>
Length		
km	kilometer	0.62 mile
m	meter	39.37 inches or 1.09 yards
cm	centimeter	0.39 inch
mm	millimeter	0.04 inch
Area		
ha	hectare	2.47 acres
Volume/Capacity		
liter	liter	61.02 cubic inches or 1.057 quarts
ml	milliliter	0.06 cubic inch or 0.034 fluid ounce
cc	cubic centimeter	0.061 cubic inch or 0.035 fluid ounce
Mass/Weight		
MT	metric ton	1.1 tons
kg	kilogram	2.205 pounds
g	gram	0.035 ounce
mg	milligram	3.5 x 10 ⁻⁵ ounce



J. Scott Angle
Dean and Director

Gerald F. Arkin
*Assistant Dean
Northern Region*

Robert N. Shulstad
*Associate Dean and
Senior Associate Director*

ISSN 0072-128X

PREFACE

This research report presents the results of the 2011 statewide performance tests of peanut, cotton and tobacco. The tests for various evaluations were conducted at several or all of the following locations: Bainbridge, Tifton, Plains and Midville in the Coastal Plain region and Athens in the Piedmont region. For identification of the test site locations, consult the map inside the back cover of this report.

Agronomic information such as grade, fiber data, plant height, lodging, disease occurrence, etc. is listed along with the yield data. Information concerning planting and harvest dates, soil type, and culture and fertilization practices used in each trial is included in footnotes.

In order to have a broad base of information, a number of varieties, including experimental lines, are included in the trials, but this does not imply that all are recommended for Georgia. Varieties best suited to a specific area or for a particular purpose and agreed upon by College of Agricultural and Environmental Sciences agronomists are presented in the 2012 Spring Planting Schedule for Georgia (available from your county Extension office). Pesticides used for production practices are included for the benefit of the reader and do not imply any endorsement or preferential treatment by the University of Georgia Agricultural Experiment Station. For additional information, contact your local county Extension agent or the nearest experiment station.

The least significant difference (LSD) at the 10% level has been included in the tables to aid in comparing hybrids. If the yields' difference of any two varieties exceeds the LSD value, they can be considered different in yield ability.

This report is one of four publications presenting the 2011 performance of agronomic crops in Georgia. For more information concerning other crops, refer to one of the following research reports: 2011 Corn Performance Tests (Annual Publication 101-3), 2010-2011 Small Grains Performance Tests (Annual Publication 100-3), 2011 Soybean, Sorghum Grain and Silage, and Summer Annual Forage Performance Tests (Annual Publication 103-3), and 2010-2011 Canola Performance data (www.swvt.uga.edu/canola).

This report, along with performance test information on other crops, is also available online at **www.swvt.uga.edu**. Additional information may be obtained by writing to J. LaDon Day, Crop and Soil Sciences Department, University of Georgia, Griffin Campus, 1109 Experiment St., Griffin, GA 30223-1797.

Cooperators

Mr. R. A. Black, Southeast Research & Education Center, Midville, Georgia
Dr. I. Flitcroft, Griffin Campus, Griffin, Georgia
Mr. S. R. Jones, Southwest Research & Education Center, Plains, Georgia
Mr. A. Knowlton, Biological Ag Engineering, Tifton Campus, Tifton, Georgia
Mr. R. R. Pines, Southwest Research & Education Center, Plains, Georgia
Mr. E. T. Ross, Tifton Campus, Tifton, Georgia
Mr. H. J. Yeomans, UGA Crop & Soil Sciences Research Farm, Athens, Georgia

Contributors

The following individuals contributed to the gathering of data and to the preparation of this report: R. Brooke, K. Cobb, J. Cox, M. Dolan, D. Dunn, M. Flynn, M. Gilmer, D. Gordan, J. Greene, J. Hudgins, R. Milton, J. Moore, J. Strickland, C. Troxell, S. Walker and G. Ware.

CONTENTS

THE SEASON with 2011 Rainfall	1
--	---

PEANUT

Tifton, Georgia:

Yield and Grade Performance, Peanut Variety Test, 2011, Irrigated.....	3
--	---

Yield and Grade Performance, Peanut Variety Test, 2011, Nonirrigated	6
--	---

Plains, Georgia:

Yield and Grade Performance, Peanut Variety Test, 2011, Irrigated.....	8
--	---

Yield and Grade Performance, Peanut Variety Test, 2011, Nonirrigated	10
--	----

Midville, Georgia:

Yield and Grade Performance, Peanut Variety Test, 2011, Irrigated.....	12
--	----

Yield and Grade Performance, Peanut Variety Test, 2011, Nonirrigated	14
--	----

COTTON

Earlier Maturity Cotton Variety Performance

Bainbridge, Georgia, 2011, Irrigated	15
--	----

Midville, Georgia, 2011, Irrigated	16
--	----

Plains, Georgia, 2011, Irrigated	17
--	----

Tifton, Georgia, 2011, Irrigated	18
--	----

Tifton, Georgia, 2011, Irrigated, including Micro-Gin Quality Data	19
--	----

Yield Summary of Earlier Maturity Cotton Varieties, 2011, Irrigated.....	20
--	----

Two-Year Summary of Earlier Maturity Cotton Varieties at Four Locations, 2010-2011, Irrigated	21
---	----

Later Maturity Cotton Variety Performance

Bainbridge, Georgia, 2011, Irrigated	22
--	----

Midville, Georgia, 2011, Irrigated	23
--	----

Plains, Georgia, 2011, Irrigated	24
--	----

Tifton, Georgia, 2011, Irrigated	25
--	----

Tifton, Georgia, 2011, Irrigated, including Micro-Gin Quality Data	26
--	----

Yield Summary of Later Maturity Cotton Varieties, 2011, Irrigated.....	27
--	----

Two-Year Summary of Later Maturity Cotton Varieties at Four Locations, 2010-2011, Irrigated	28
---	----

Cotton Strains Performance

Midville, Georgia, 2011, Irrigated	29
--	----

Plains, Georgia, 2011, Irrigated	30
--	----

Tifton, Georgia, 2011, Irrigated	31
--	----

Yield Summary of Cotton Strains, 2011, Irrigated.....	32
---	----

Dryland Earlier Maturity Cotton Variety Performance

Athens, Georgia, 2011 - Earlier Maturity.....	33
---	----

Midville, Georgia, 2011 - Earlier Maturity.....	34
---	----

Plains, Georgia, 2011 - Earlier Maturity	35
--	----

Tifton, Georgia, 2011 - Earlier Maturity.....	36
---	----

Yield Summary of Dryland Earlier Maturity Cotton Varieties, 2011	37
--	----

Two-Year Summary of Dryland Earlier Maturity Cotton Varieties at Four Locations, 2010-2011	38
--	----

Dryland Later Maturity Cotton Variety Performance

Athens, Georgia, 2011 - Later Maturity.....	39
---	----

Midville, Georgia, 2011 - Later Maturity	40
--	----

Plains, Georgia, 2011 - Later Maturity	41
--	----

Tifton, Georgia, 2011 - Later Maturity	42
--	----

Yield Summary of Dryland Later Maturity Cotton Varieties, 2011.....	43
---	----

Two-Year Summary of Dryland Later Maturity Cotton Varieties at Four Locations, 2010-2011	44
--	----

TOBACCO

Tifton, Georgia:

Official Flue-Cured Tobacco Variety Test - Yield, Value, Price Index, Grade Index, and	
--	--

Agronomic and Chemical Characteristics of Released Varieties, 2011	45
--	----

Three- and Two-Year Averages of Official Flue-Cured Tobacco Variety Tests -	
---	--

Comparison of Released Varieties for Certain Characteristics, 2008, 2010 and 2011	46
---	----

Regional Farm Flue-Cured Tobacco Variety Test - Comparison of Released Varieties	
--	--

for Certain Characteristics, 2011.....	48
--	----

2011 PEANUT, COTTON AND TOBACCO PERFORMANCE TESTS

*J. LaDon Day, Anton E. Coy, Stevan S. LaHue,
Larry G. Thompson and John D. Gassett, Editors*

The Season

The spring of 2011 began with abnormally dry soil, completely different than the March 2010 wet and cold soil conditions. However, during early spring most areas did have enough moisture for seeding. Planting progressed ahead of five-year averages. Plant stands and early season growth were good in most areas. Due to a dry April, in early May less than half of the state had adequate moisture, and as the lack of rainfall and high temperatures continued into the end of May, two-thirds of the state was under a severe drought. Producers quickly fell behind in crop progress to late planting their crops or not planting at all. Most of the non-irrigated crops were severely damaged beyond salvage from the high heat and lack of moisture. Irrigation, which began at planting, struggled to keep up over much of the state throughout the summer and fall. Insects were a concern in most areas. Diseases, especially white mold, developed into an issue for the majority of the peanut growing area of the state and persisted throughout the growing season.

Rainfall amounts recorded monthly at the five test locations in Georgia during the 2011 growing season are presented in the following table. Seasonal rainfall totals were below normal all across the state. The average rainfall deficit was near 12 inches or 34% in the Piedmont and Coastal Plain regions, with the only exception being the Tifton area where 82% of normal rainfall occurred. Extremely dry conditions (60% of normal rainfall) persisted for the last two years in Burke (Midville) and Sumter (Plains) counties and surrounding areas.

2011 Rainfall¹

Month	Athens ²	Attapulgus ³	Midville	Plains	Tifton
March	7.89	4.51	2.85	3.02	1.54
April	2.50	2.36	1.41	2.01	1.60
May	1.40	1.05	2.33	0.03	0.01
June	3.69	4.84	1.73	1.12	4.48
July	1.90	5.68	3.15	7.00	6.26
August	1.00	1.55	1.02	1.50	1.82
September	1.63	3.31	2.40	3.38	5.71
October	3.57	4.21	2.46	1.22	4.81
November	2.78	3.27	2.05	1.96	1.34
Total	26.36	30.78	19.40	21.24	27.57
Normal (9 mo)*	35.92	41.76	32.60	35.23	33.65

1. Data provided in part by Dr. I. Flitcroft, Georgia Station, Griffin, GA.

2. Plant Sciences Farm.

3. Attapulgus Research Center is nearest location to the Bainbridge site.

J. LaDon Day is the program director of the statewide variety testing program and John D. Gassett is a research professional II in the Department of Crop and Soil Sciences, Griffin Campus, Griffin, GA 30223-1797. Anton E. Coy, Stevan S. LaHue and Larry G. Thompson are senior agricultural specialist, agricultural specialist and research professional I, respectively, in the Department of Crop and Soil Sciences, Tifton Campus, Tifton, GA 31793-0748.

Crop maturity progressed ahead of the five-year average and harvest conditions during 2011 were excellent. Georgia peanut producers planted 475,000 acres this year, a 16% decrease from last year and the lowest planting in 29 years. During 2011 cotton farmers seeded 1.60 million acres, 20% more than last year. Tobacco planting in the state abated, reflecting a steady 10-year decline as 11,500 acres were transplanted during the 2011 crop year.

The state per acre yield for peanuts was 3,400 pounds, which matched 2008 but produced 18% less than last year and the lowest tonnage of nuts produced in nine years. Cotton per acre yield in 2011 of 837 pounds was 2% higher than last year and the highest per acre yield in six years. This yield level totaled over harvested acres of cotton produced a new record for cotton production in Georgia. Tobacco production in the state of Georgia continued to decline due to a 2% decrease in per-acre yield during 2011 compared to 2010.

PEANUT

Tifton, Georgia: Yield and Grade Performance Peanut Variety Trial, 2011, Irrigated

Variety	Digging Date	Yield lb/A	TSMK %	OK %	DK %	ELK %	Seed no./lb	Fancy %
<u>Spanish Types</u>								
GA 082549 ¹	09/30	4870	77.0	3.0	0.0	0.0	868	0.0
Georgia-04S	09/30	4795	74.5	4.0	0.5	0.0	1015	0.0
GA 082550 ¹	10/17	4779	75.5	3.5	0.5	0.0	1066	0.0
GA 082548 ¹	09/30	4764	75.5	3.5	0.0	0.0	966	0.0
GA 082554 ¹	10/17	4731	76.5	3.0	0.5	0.0	1054	0.0
GA 082551 ¹	10/17	4577	78.0	3.0	0.0	0.0	1079	0.0
GA 082552 ¹	10/17	4559	76.0	4.5	0.0	0.0	1001	0.0
GA 082553 ¹	10/17	4426	76.5	3.5	0.0	0.0	937	0.0
Georgia Browne	09/30	3966	72.5	6.0	0.0	0.0	1045	0.0
Tamspan 90	08/23	2883	62.0	9.0	0.0	0.0	1127	0.0
Tamnut OL06	08/23	2862	62.0	5.5	0.5	0.0	905	0.0
OLin	08/23	2333	64.0	6.5	0.5	0.0	1186	0.0
Pronto	08/12	2263	65.0	8.0	0.5	0.0	1182	0.0
Spanco	08/12	2217	63.0	7.0	1.0	0.0	1145	0.0
Average	09/21	3859	71.3	5.0	0.3	0.0	1041	0.0
LSD at 10% Level		494	3.3	2.4	-	-	98	-
C.V. %		16.1	2.9	20.0	-	-	5.0	-
<u>Valencia Types</u>								
Valencia McRan	08/12	1717	56.5	10.5	0.5	0.0	1173	0.0
Georgia Red	08/23	1716	61.5	8.5	0.5	0.0	966	0.0
N.M. Valencia A	08/12	1690	54.5	11.5	0.5	0.0	1165	0.0
N.M. Valencia C	08/12	1667	52.5	13.0	0.5	0.0	1176	0.0
H & W Valencia 136	08/12	1582	49.5	14.0	0.5	0.0	1241	0.0
Georgia Valencia	08/23	1558	48.5	8.0	5.0	0.0	910	0.0
Average	08/16	1655	53.8	10.9	1.3	0.0	1105	0.0
LSD at 10% Level		494	3.3	2.4	-	-	98	-
C.V. %		16.1	2.9	20.0	-	-	5.0	-

1. Advanced Georgia breeding line.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 9, 2011.

Seeding Rate: 6 seed/row foot in 36" rows.

Fertilization: 0 lb N, 0 lb P₂O₅, and 0 lb K₂O/acre.

Soil Test: P = High, K = High, and pH = 6.5.

Soil Type: Tifton sandy loam.

Previous Crop: Cotton.

Management: Disked, moldboard plowed and rototilled; Sonalan, Dual Magnum, Ultra Blazer and Basagran used for weed control; Provost, Folicur, Clorothalnil and Artisan used for fungal control; 1000 lb/acre Landplaster; irrigated 9 inches.

Test conducted by A. Coy, R. Brooke and D. Dunn.

Tifton, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Irrigated

Variety	Digging Date	Yield lb/A	TSMK %	OK %	DK %	ELK %	Seed no./lb	Fancy %
<u>Runner Types</u>								
GA 072531 ¹	10/17	6434	73.5	3.5	0.5	0.0	723	0.0
GA 082546 ¹	10/17	6089	77.0	3.5	0.0	0.0	891	0.0
GA 072716 ¹	09/30	6002	75.5	3.5	0.0	0.0	792	0.0
FloRun™ '107'	09/30	5433	73.5	4.0	0.0	0.0	712	0.0
GA 082524 ¹	10/17	5421	75.5	4.0	0.5	0.0	923	0.0
GA 072515 ¹	09/16	5385	79.0	2.0	0.5	0.0	693	0.0
Georgia-06G	09/16	5345	77.0	2.0	0.5	0.0	623	0.0
GA 072514 ¹	09/16	5276	82.0	1.0	0.0	0.0	705	0.0
Georgia-10T	10/17	5273	78.5	2.5	0.5	0.0	669	0.0
Florida-07	09/30	5260	72.5	2.5	0.0	0.0	585	0.0
Georgia Greener	09/16	5185	77.5	2.5	0.0	0.0	708	0.0
Georgia-07W	09/30	5115	76.0	2.5	0.5	0.0	688	0.0
GA 082522 ¹	09/30	5106	75.0	5.0	0.0	0.0	828	0.0
Tifguard	09/16	5061	75.0	2.0	0.0	0.0	617	0.0
GA 072523 ¹	09/16	4991	78.5	1.5	0.0	0.0	648	0.0
UF 10302 ²	09/16	4982	76.0	1.5	1.5	0.0	653	0.0
Georgia-09B	09/16	4949	77.0	2.0	0.0	0.0	721	0.0
Georgia-02C	09/30	4604	74.5	4.0	0.0	0.0	835	0.0
GA 082549 ¹	09/30	4586	76.0	3.0	0.0	0.0	909	0.0
GA 082548 ¹	09/30	4547	74.0	5.0	0.0	0.0	962	0.0
Georgia Green	09/16	4483	76.0	3.0	0.5	0.0	813	0.0
Average	09/27	5216	76.2	2.9	0.2	0.0	747	0.0
LSD at 10% Level		508	3.2	-	-	-	47	-
C.V. %		10.4	2.6	-	-	-	4.1	-
<u>Virginia Types</u>								
Georgia-08V	09/16	5376	71.5	2.0	3.0	61.5	466	89.5
CHAMPS	08/23	5212	72.5	1.5	0.0	42.5	478	86.5
Bailey	08/23	5200	69.0	3.5	0.0	40.0	508	86.5
Georgia-11J	10/17	5161	74.5	1.0	0.5	60.5	406	83.5
Gregory	08/23	4925	68.0	2.0	0.0	44.5	516	86.5
Sugg	08/23	4895	70.5	2.0	0.0	48.5	499	85.5
Florida Fancy	09/16	4668	69.5	1.5	0.0	45.0	500	84.5
Titan	08/23	4356	63.0	1.5	0.5	30.5	473	93.0
Perry	08/23	4280	68.5	3.5	0.0	33.5	551	76.0
Average	09/03	4897	69.7	2.1	0.4	45.2	488	85.7
LSD at 10% Level		508	3.2	-	-	5.7	47	5.7
C.V. %		10.4	2.6	-	-	24.7	4.1	10

Tifton, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Irrigated (Continued)

1. Advanced Georgia breeding line.
2. Advanced Florida breeding line.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD ($P = 0.10$).

Planted: May 9, 2011.
Seeding Rate: 6 seed/row foot in 36" rows.
Fertilization: 0 lb N, 0 lb P_2O_5 , and 0 lb K_2O /acre.
Soil Test: P = High, K = High, and pH = 6.5.
Soil Type: Tifton sandy loam.
Previous Crop: Cotton.
Management: Disked, moldboard plowed and rototilled; Sonalan, Dual Magnum, Ultra Blazer and Basagran used for weed control; Provost, Folicur, Clorothilanil and Artisan used for fungal control; 1000 lb/acre Landplaster; irrigated 9 inches.

Test conducted by A. Coy, R. Brooke and D. Dunn.

Tifton, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Nonirrigated

Variety	Digging Date	Yield lb/A	TSMK %	OK %	DK %	ELK %	Seed no./lb	Fancy %
<u>Runner Types</u>								
Florida-07	10/28	5364	66.5	4.0	2.0	0.0	591	0.0
GA 072531 ¹	10/28	5282	70.0	5.5	0.5	0.0	735	0.0
Georgia Greener	10/17	5179	70.5	5.5	1.0	0.0	731	0.0
Georgia-07W	10/28	4924	73.5	3.0	0.5	0.0	642	0.0
Georgia-10T	10/28	4725	69.0	6.0	3.0	0.0	700	0.0
Georgia-06G	10/17	4580	73.0	4.0	0.5	0.0	772	0.0
GA 072523 ¹	10/17	4559	72.0	3.5	0.5	0.0	788	0.0
GA 082549 ¹	10/28	4520	71.0	6.5	0.5	0.0	877	0.0
GA 072716 ¹	10/28	4429	62.5	10.0	2.0	0.0	941	0.0
GA 082522 ¹	10/28	4413	72.0	6.0	1.5	0.0	793	0.0
FloRun™ '107'	10/28	4383	68.5	6.5	1.5	0.0	738	0.0
Tifguard	10/17	4350	69.0	5.5	0.5	0.0	677	0.0
GA 072515 ¹	10/17	4292	72.5	4.0	0.5	0.0	949	0.0
GA 082546 ¹	10/28	4274	68.5	8.0	0.5	0.0	867	0.0
UF 10302 ²	10/17	4117	69.5	5.5	0.0	0.0	678	0.0
Georgia-09B	10/17	4029	68.0	7.0	0.5	0.0	777	0.0
Georgia Green	10/17	4013	69.0	6.5	1.0	0.0	833	0.0
GA 082548 ¹	10/28	4005	71.0	5.0	1.0	0.0	926	0.0
Georgia-02C	10/28	4002	71.0	4.5	1.0	0.0	937	0.0
GA 072514 ¹	10/17	3933	75.5	3.5	0.5	0.0	933	0.0
GA 082524 ¹	10/28	3442	67.0	8.5	1.5	0.0	925	0.0
Average	10/23	4420	70.0	5.6	1.0	0.0	800	0.0
LSD at 10% Level		501	6.1	-	-	-	109	-
C.V. %		12.6	5.3	-	-	-	8.9	-
<u>Virginia Types</u>								
Georgia-11J	10/28	4779	64.5	3.0	0.5	43.5	523	74.5
Georgia-08V	10/17	4244	63.0	4.5	4.0	39.5	522	75.0
CHAMPS	09/16	3830	65.5	3.0	1.5	39.5	515	88.5
Gregory	09/16	3805	61.0	3.0	1.0	32.0	575	80.0
Bailey	09/16	3748	69.0	1.5	0.0	40.0	504	79.5
Sugg	09/16	3213	66.0	2.5	0.0	41.5	525	82.5
Florida Fancy	10/17	3198	58.5	5.0	2.0	25.5	566	72.0
Titan	09/16	3110	53.0	4.5	1.0	26.5	585	82.0
Perry	09/16	2287	67.5	2.5	1.0	39.0	553	69.5
Average	09/28	3579	63.1	3.3	1.2	36.3	541	78.2
LSD at 10% Level		501	6.1	-	-	6.5	109	6.5
C.V. %		12.6	5.3	-	-	3.8	8.9	16.3

Tifton, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Nonirrigated (Continued)

1. Advanced Georgia breeding line.
2. Advanced USDA breeding line.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD ($P = 0.10$).

Planted: May 10, 2011.
Seeding Rate: 6 seed/row foot in 36" rows.
Fertilization: 0 lb N, 0 lb P_2O_5 , and 0 lb K_2O /acre.
Soil Test: P = High, K = High, and pH = 5.8.
Soil Type: Tifton loamy sand.
Previous Crop: Corn and Grain Sorghum.
Management: Disked, moldboard plowed and rototilled; Sonalan, Dual Magnum, Basagran, 24DB and Select used for weed control; Provost, Folicur, Clorothilanil and Artisan used for fungal control; 1000 lb/acre Landplaster.

Test conducted by A. Coy, R. Brooke and D. Dunn.

Plains, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Irrigated

Variety	Digging Date	Yield lb/A	TSMK %	OK %	DK %	ELK %	Seed no./lb	Fancy %
<u>Runner Types</u>								
GA 072716 ¹	10/05	6271	73.0	4.5	0.0	0.0	852	0.0
Georgia-06G	10/05	6207	75.0	3.0	0.5	0.0	798	0.0
GA 082522 ¹	10/05	5729	74.0	4.5	1.5	0.0	876	0.0
Georgia-07W	10/05	5554	76.0	3.0	0.0	0.0	658	0.0
FloRun™ '107'	10/05	5492	72.5	4.0	0.5	0.0	788	0.0
Georgia-02C	10/05	5469	74.5	3.5	0.0	0.0	776	0.0
Georgia-09B	10/05	5333	76.0	3.0	0.5	0.0	747	0.0
Georgia Greener	10/05	5236	77.0	4.5	0.0	0.0	725	0.0
Florida-07	10/05	5218	71.0	3.5	0.5	0.0	730	0.0
UF 10302 ²	10/05	5170	73.0	5.0	0.0	0.0	654	0.0
GA 072531 ¹	10/26	5106	72.0	3.5	0.0	0.0	788	0.0
GA 072523 ¹	10/05	5049	76.0	2.5	0.0	0.0	702	0.0
GA 072515 ¹	10/05	5049	78.5	2.0	0.0	0.0	767	0.0
Georgia Green	10/05	4893	75.5	3.5	0.0	0.0	731	0.0
GA 072514 ¹	10/05	4770	78.5	2.5	0.5	0.0	744	0.0
GA 082548 ¹	10/05	4625	75.0	4.0	0.0	0.0	916	0.0
Tifguard	10/05	4532	73.5	3.5	0.5	0.0	724	0.0
GA 082546 ¹	10/26	4513	73.5	5.5	0.0	0.0	935	0.0
GA 082524 ¹	10/26	4389	77.0	3.5	0.0	0.0	831	0.0
GA 082549 ¹	10/05	4356	76.0	3.0	0.0	0.0	961	0.0
Georgia-10T	10/26	4127	77.0	3.5	0.0	0.0	655	0.0
Average	10/09	5099	75.0	3.6	0.2	0.0	779	0.0
LSD at 10% Level		411	3.5	-	-	-	12.1	-
C.V. %		8.6	2.8	-	-	-	10.5	-
<u>Virginia Types</u>								
Georgia-08V	10/05	5757	75.0	1.0	0.5	61.0	404	86.5
Bailey	09/26	4907	72.0	1.5	0.0	47.5	501	85.5
Gregory	09/26	4819	66.5	3.5	0.0	38.5	447	89.5
Florida Fancy	10/05	4734	73.0	1.0	0.0	57.0	422	92.5
CHAMPS	09/26	4692	68.5	1.5	0.0	53.0	427	92.0
Sugg	09/26	4665	72.5	1.5	0.0	51.5	477	87.0
Perry	09/26	4365	71.5	2.0	1.0	48.0	477	77.0
Georgia-11J	10/26	4309	75.0	1.0	0.0	62.0	409	89.0
Titan	09/26	4105	69.0	2.5	1.5	48.0	436	93.0
Average	10/01	4706	71.4	1.7	0.3	51.8	444	88.0
LSD at 10% Level		411	3.5	-	-	5.3	12.1	2.2
C.V. %		8.6	2.8	-	-	20	10.5	5

**Plains, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Irrigated (Continued)**

1. Advanced Georgia breeding line.
2. Advanced USDA breeding line.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD ($P = 0.10$).

Planted: May 19, 2011.
Seeding Rate: 6 seed/row foot in 36" rows.
Fertilization: 0 lb N, 0 lb P_2O_5 , and 0 lb K_2O /acre.
Soil Test: P = High, K = Very High, and pH = 5.9.
Soil Type: Greenville loamy sand.
Previous Crop: Corn.
Management: Disked, moldboard plowed and rototilled; Valor, Strongarm and Sonalan used for weed control; Lorsban used for insect control; Provost used for fungal control; 1000 lb/acre lime; irrigated 10 inches.

Test conducted by A. Coy, R. Brooke, D. Dunn and R. Pines.

Plains, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Nonirrigated

Variety	Digging Date	Yield	TSMK	OK	DK	ELK	Seed	Fancy
		lb/A	%	%	%	%	no./lb	%
Runner Types								
GA 072716 ¹	10/26	3987	68.5	7.0	0.0	0.0	910	0.0
Georgia-07W	10/26	3910	70.5	4.0	0.0	0.0	764	0.0
Georgia-06G	10/26	3815	70.5	4.5	0.0	0.0	683	0.0
Georgia-09B	10/26	3582	71.5	4.5	0.0	0.0	762	0.0
FloRun™ '107'	10/26	3570	69.5	6.0	0.0	0.0	904	0.0
GA 082522 ¹	10/26	3554	74.0	5.0	0.0	0.0	833	0.0
GA 072531 ¹	10/26	3503	65.5	7.0	0.0	0.0	781	0.0
GA 082549 ¹	10/26	3503	72.5	4.0	0.0	0.0	993	0.0
Florida-07	10/26	3481	67.0	6.0	0.0	0.0	747	0.0
Georgia-02C	10/26	3458	69.0	5.5	0.0	0.0	820	0.0
GA 072523 ¹	10/26	3395	74.5	2.0	0.0	0.0	759	0.0
GA 072515 ¹	10/26	3358	73.5	3.0	0.0	0.0	760	0.0
Georgia Greener	10/26	3307	73.5	3.5	0.5	0.0	758	0.0
GA 072514 ¹	10/26	3260	73.0	5.0	0.0	0.0	785	0.0
UF 10302 ²	10/26	3246	71.5	3.5	0.0	0.0	712	0.0
Tifguard	10/26	3140	70.5	4.5	0.0	0.0	688	0.0
GA 082548 ¹	10/26	3140	71.5	5.0	0.0	0.0	1070	0.0
GA 082524 ¹	10/26	3010	72.0	5.0	0.0	0.0	879	0.0
GA 082546 ¹	10/26	2983	71.0	5.5	0.0	0.0	748	0.0
Georgia Green	10/26	2964	69.0	7.0	0.0	0.0	839	0.0
Georgia-10T	10/26	2849	73.5	4.5	0.0	0.0	726	0.0
Average	10/26	3382	71.0	4.9	0.0	0.0	806	0.0
LSD at 10% Level		439	2.7	-	-	-	98	-
C.V. %		13.8	2.2	-	-	-	8.0	-
Virginia Types								
Georgia-08V	10/26	3673	72.5	1.5	0.0	45.5	499	74.5
Sugg	10/05	3639	70.5	3.0	0.5	47.5	471	76.0
Gregory	10/05	3582	67.5	2.5	1.0	51.0	488	87.0
CHAMPS	10/05	3518	68.5	2.5	0.5	34.0	478	83.5
Bailey	10/05	3319	66.5	4.0	0.0	19.5	584	69.0
Georgia-11J	10/26	2955	68.5	2.5	0.5	43.5	535	64.0
Perry	10/05	2929	71.0	3.0	0.0	33.5	529	60.5
Titan	10/05	2716	62.0	2.0	1.0	41.5	454	90.5
Florida Fancy	10/26	2384	67.0	3.0	0.0	28.5	575	72.5
Average	10/12	3190	68.2	2.7	0.4	38.3	512	75.3
LSD at 10% Level		439	2.7	-	-	4.1	98	6.7
C.V. %		13.8	2.2	-	-	20.9	8.0	17.4

Plains, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Nonirrigated (Continued)

1. Advanced Georgia breeding line.
2. Advanced USDA breeding line.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD ($P = 0.10$).

Planted: May 23, 2011.
Seeding Rate: 6 seed/row foot in 36" rows.
Fertilization: 0 lb N, 0 lb P_2O_5 , and 0 lb K_2O /acre.
Soil Test: P = High, K = Very High, and pH = 5.9.
Soil Type: Greenville loamy sand.
Previous Crop: Cotton.
Management: Disked, moldboard plowed and rototilled; Valor, Strongarm and Sonalan used for weed control; Lorsban used for insect control; Provost used for fungal control.

Test conducted by A. Coy, R. Brooke, D. Dunn and R. Pines.

Midville, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Irrigated

Variety	Digging Date	Yield	TSMK	OK	DK	ELK	Seed	Fancy
		lb/A	%	%	%	%	no./lb	%
<u>Runner Types</u>								
GA 072716 ¹	10/17	7556	77.0	3.5	0.0	0.0	812	0.0
GA 072531 ¹	10/17	7458	76.0	2.0	0.0	0.0	749	0.0
Georgia-07W	10/17	7161	77.5	3.0	0.0	0.0	658	0.0
FloRun™ '107'	10/17	6912	76.5	2.5	0.0	0.0	674	0.0
Georgia-09B	10/01	6888	76.5	2.0	0.0	0.0	648	0.0
GA 072515 ¹	10/01	6864	78.0	2.5	0.0	0.0	694	0.0
Georgia-06G	10/01	6814	77.0	2.0	0.5	0.0	592	0.0
GA 082524 ¹	10/17	6785	77.0	2.5	0.0	0.0	804	0.0
Tifguard	10/01	6768	74.5	3.0	0.0	0.0	606	0.0
GA 082522 ¹	10/17	6767	78.0	4.0	0.0	0.0	799	0.0
Florida-07	10/17	6707	74.0	3.0	0.0	0.0	628	0.0
Georgia Greener	10/01	6590	77.0	1.5	0.0	0.0	663	0.0
GA 082546 ¹	10/17	6583	76.5	2.5	0.0	0.0	838	0.0
Georgia-10T	10/17	6458	80.5	1.5	0.0	0.0	671	0.0
GA 082549 ¹	10/17	6367	77.0	2.5	0.0	0.0	808	0.0
GA 072514 ¹	10/01	6357	79.5	2.5	0.0	0.0	713	0.0
GA 072523 ¹	10/01	6248	78.5	1.5	0.0	0.0	677	0.0
UF 10302 ²	10/01	6189	76.0	3.0	0.0	0.0	649	0.0
GA 082548 ¹	10/17	6088	75.5	4.5	0.0	0.0	943	0.0
Georgia-02C	10/17	5994	75.0	3.0	0.5	0.0	766	0.0
Georgia Green	10/01	5919	76.0	3.5	0.0	0.0	796	0.0
Average	10/10	6642	76.8	2.7	0.0	0.0	723	0.0
LSD at 10% Level		584	2.5	-	-	-	57	-
C.V. %		9.7	2.0	-	-	-	5.2	-
<u>Virginia Types</u>								
Georgia-11J	10/17	7239	76.5	1.0	0.0	60.0	412	85.0
Georgia-08V	10/01	6637	75.0	1.5	0.5	56.0	504	71.0
Florida Fancy	10/01	6014	72.0	2.0	0.5	47.0	495	73.0
Gregory	09/19	5285	70.5	1.5	0.5	58.0	448	90.0
Sugg	09/19	4851	71.5	3.0	0.0	43.5	495	82.0
Bailey	09/19	4834	72.0	3.0	0.5	48.0	500	84.5
CHAMPS	09/19	4795	73.5	1.5	0.5	48.0	468	84.5
Titan	09/19	4784	69.0	1.0	0.5	49.5	420	88.0
Perry	09/19	4637	71.5	2.5	0.0	37.5	503	80.5
Average	09/25	5453	72.4	1.9	0.3	49.7	472	82.1
LSD at 10% Level		584	2.5	-	-	3.6	57	2.7
C.V. %		9.7	2.0	-	-	14.2	5.2	6.4

Midville, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Irrigated (Continued)

1. Advanced Georgia breeding line.
2. Advanced USDA breeding line.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD ($P = 0.10$).

Planted: May 12, 2011.
Seeding Rate: 6 seed/row foot in 36" rows.
Fertilization: 0 lb N, 0 lb P_2O_5 , and 0 lb K_2O /acre.
Soil Test: P = High, K = Very High, and pH = 5.8.
Soil Type: Tifton sandy loam.
Previous Crop: Corn.
Management: Disked, moldboard plowed and field conditioned; Sonalan, Valor, Acumen, Gramoxone, Storm and Duel used for weed control; Headline, Folicur, Convoy and Chlorothalonil used for fungal control; 1000 lb/a Dolemite Lime and 1000 lb/a Gypsum; irrigated 15 inches.

Test conducted by A. Coy, R. Brooke, D. Dunn, K. Cobb and R. Milton.

Midville, Georgia:
Yield and Grade Performance
Peanut Variety Trial, 2011, Nonirrigated

Peanut varieties were planted at this location on May 12, 2011 in a nonirrigated test. However, extensive damage throughout the growing season from lack of rainfall, high temperatures and droughty growing conditions resulted in a peanut crop not useful for performance evaluation.

Test conducted by A. Coy, R. Brooke, D. Dunn, K. Cobb and R. Milton.

COTTON

Bainbridge, Georgia: Earlier Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
DP 1028 B2RF	1942	46.8	84.7	1.14	28.4	5.1
DP 1133 B2RF	1928	46.5	85.0	1.16	31.7	5.0
AM1511 B2RF	1886	45.1	84.2	1.15	31.1	4.5
PHY 499 WRF	1877	46.4	84.9	1.14	31.2	5.0
DP 0912 B2RF	1847	43.3	83.4	1.13	29.6	5.1
CG 3787 B2RF	1781	45.7	85.0	1.16	28.5	4.9
Dyna-Gro 2570B2RF	1761	43.4	84.9	1.15	29.2	4.8
BRS286	1707	42.4	84.0	1.14	31.8	4.6
GA2006106	1691	42.0	85.0	1.19	33.9	4.8
PHY 375 WRF	1684	44.6	83.9	1.15	29.0	4.6
BX 1252LLB2	1681	42.1	83.9	1.17	30.9	4.9
BX 1262B2F	1667	43.3	83.9	1.17	29.2	4.8
DP 0949B2RF	1655	44.6	84.5	1.15	30.4	5.0
PHY 367 WRF	1640	43.8	84.6	1.19	29.6	4.5
BCSX 1150B2RF	1626	39.6	85.4	1.22	32.9	4.7
GA2004143	1617	44.5	84.8	1.18	34.9	4.9
ST 4288B2F	1606	40.3	83.5	1.15	26.9	4.8
All-Tex ATX3039 B2RF	1593	43.7	84.0	1.14	29.1	4.5
FM1740B2RF	1580	42.9	83.7	1.16	29.8	4.7
ST 4145LLB2	1566	42.7	84.8	1.17	33.7	4.5
BRS293	1542	42.0	84.5	1.17	33.4	4.9
AM 1550 B2RF	1526	42.5	83.5	1.14	27.1	4.9
All-Tex LA122	1524	43.8	85.1	1.17	29.8	4.6
SSG CT310 HQ	1522	41.3	83.8	1.15	33.3	4.8
DP 0920 B2RF	1507	43.8	83.9	1.15	28.3	4.8
All-Tex 7A21	1503	44.2	84.3	1.17	30.2	4.8
All-Tex ATX81144	1481	42.7	85.8	1.25	31.9	4.0
DP 0924 B2RF	1468	41.9	85.1	1.15	30.7	5.1
SSG HQ 210 CT	1399	40.6	82.3	1.13	30.3	4.7
GA2008057	1302	40.0	84.8	1.19	33.9	4.4
SSG CT Linwood	1185	41.4	84.3	1.13	33.8	5.2
Average	1622	43.2	84.3	1.16	30.8	4.7
LSD 0.10	190	0.8	1.2	0.03	2.3	0.3
CV %	10.0	1.6	0.8	1.82	4.4	4.2

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 11, 2011.

Harvested: October 17, 2011.

Soil Type: Bonnaeu blanton loamy sand.

Fertilization: 180 lb N, 71 lb P₂O₅, and 122 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	May	June	July	Aug.	Sept.
Irrigation (in):	1.60	4.80	1.60	3.20	0.80
Rainfall (in):	2.10	1.35	9.10	2.35	5.05

Trials conducted by Larry Thompson.

Midville, Georgia: Earlier Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
PHY 499 WRF	2957	47.7	84.5	1.14	31.1	4.7
DP 0912 B2RF	2924	45.8	83.3	1.12	27.8	4.6
DP 1028 B2RF	2777	46.1	85.2	1.18	28.3	5.0
All-Tex ATX3039 B2RF	2741	45.2	84.6	1.16	28.8	4.6
AM1511 B2RF	2637	47.9	84.8	1.16	29.6	5.1
DP 0949B2RF	2610	44.2	84.9	1.18	30.0	4.8
DP 0924 B2RF	2599	44.4	84.4	1.15	30.2	5.0
GA2004143	2597	44.7	84.8	1.22	32.6	4.6
BRS293	2582	44.2	84.4	1.16	31.3	4.8
BCSX 1150B2RF	2580	42.1	84.5	1.19	32.5	5.2
BX 1252LLB2	2567	43.7	83.7	1.16	30.9	4.6
CG 3787 B2RF	2547	43.6	84.2	1.16	30.4	4.6
GA2006106	2534	43.2	85.2	1.25	33.2	4.7
BX 1262B2F	2534	44.8	84.7	1.16	31.3	5.1
FM1740B2RF	2532	44.8	83.4	1.12	27.8	5.0
All-Tex ATX81144	2530	42.8	83.7	1.23	32.1	4.2
PHY 375 WRF	2528	44.7	84.4	1.16	28.7	4.7
DP 0920 B2RF	2510	45.5	83.9	1.15	26.9	4.8
ST 4145LLB2	2494	43.4	85.4	1.18	31.2	4.6
All-Tex 7A21	2490	44.7	84.8	1.19	30.2	4.6
SSG HQ 210 CT	2451	41.2	83.4	1.16	30.5	4.8
AM 1550 B2RF	2439	42.7	84.5	1.13	28.1	4.8
Dyna-Gro 2570B2RF	2426	42.0	84.1	1.17	29.8	4.8
All-Tex LA122	2387	45.6	84.9	1.17	28.5	4.3
SSG CT Linwood	2347	43.9	85.1	1.13	33.7	5.4
ST 4288B2F	2341	42.2	83.9	1.17	27.0	4.9
DP 1133 B2RF	2335	47.0	85.6	1.18	31.9	4.8
PHY 367 WRF	2328	43.6	84.4	1.15	28.9	4.7
SSG CT310 HQ	2197	41.4	84.7	1.16	33.3	4.9
BRS286	2070	42.1	83.1	1.11	31.8	4.8
GA2008057	1750	42.0	85.5	1.21	32.2	4.3
Average	2495	44.1	84.4	1.16	30.3	4.7
LSD 0.10	244	1.5	N.S. ¹	0.05	2.8	0.5
CV %	8.4	2.8	1.0	2.47	5.4	6.0

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 3, 2011.

Harvested: October 3, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 96 lb N, 80 lb P₂O₅, and 60 lb K₂O/acre.

Management: Temik applied 5 lb/acre and Telone II applied 3 gal/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0	1.90	2.45	4.00	4.00	0
Rainfall (in):	1.41	2.32	1.76	2.22	1.50	2.51

Trials conducted by Larry Thompson.

Plains, Georgia:
Earlier Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
ST 4145LLB2	2150	41.6	83.9	1.20	31.2	4.0
DP 0912 B2RF	2022	42.6	84.9	1.17	30.3	4.5
AM1511 B2RF	1957	44.1	84.7	1.18	29.4	4.5
DP 1133 B2RF	1947	44.1	85.1	1.21	30.9	4.3
PHY 367 WRF	1931	42.1	84.9	1.21	29.9	4.1
ST 4288B2F	1924	39.2	84.6	1.22	31.1	4.2
All-Tex ATX3039 B2RF	1912	43.0	84.0	1.19	28.4	3.9
CG 3787 B2RF	1908	43.9	85.3	1.18	28.2	4.4
BX 1252LLB2	1849	40.8	85.3	1.22	30.7	3.9
All-Tex 7A21	1848	43.0	85.4	1.23	32.2	4.1
BX 1262B2F	1821	41.7	85.7	1.21	31.2	4.3
PHY 375 WRF	1806	43.0	84.4	1.19	28.3	4.0
DP 0920 B2RF	1798	42.0	85.2	1.18	28.9	4.4
PHY 499 WRF	1778	45.7	85.1	1.19	32.8	4.3
All-Tex LA122	1772	43.7	85.2	1.21	29.4	4.3
AM 1550 B2RF	1752	41.2	84.6	1.18	28.9	4.3
Dyna-Gro 2570B2RF	1722	42.4	85.7	1.20	31.7	4.0
BCSX 1150B2RF	1677	38.0	85.6	1.26	34.6	4.3
FM1740B2RF	1667	41.3	84.6	1.22	29.4	3.9
DP 1028 B2RF	1664	44.6	85.4	1.17	29.0	4.6
DP 0924 B2RF	1661	41.5	84.5	1.16	30.2	4.2
DP 0949B2RF	1624	42.9	84.8	1.20	30.4	4.4
All-Tex ATX81144	1595	40.3	85.6	1.27	32.2	3.5
GA2004143	1577	43.2	86.1	1.26	33.3	4.0
BRS286	1567	40.6	83.8	1.18	31.6	4.2
SSG HQ 210 CT	1541	40.6	84.3	1.17	32.9	4.3
GA2008057	1452	40.9	85.3	1.23	34.4	4.0
GA2006106	1358	40.5	85.0	1.25	33.7	3.8
SSG CT Linwood	1256	41.6	84.9	1.15	33.1	4.6
BRS293	1246	39.2	83.6	1.20	37.5	4.3
SSG CT310 HQ	1194	39.1	84.7	1.18	34.7	4.5
Average	1709	41.9	84.9	1.20	31.3	4.2
LSD 0.10	270	1.0	1.0	0.02	2.5	0.3
CV %	13.4	2.1	0.7	1.42	4.7	4.1

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 5, 2011.

Harvested: October 7, 2011.

Soil Type: Greenville sandy clay loam.

Fertilization: 100 lb N, 88 lb P₂O₅, and 24 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0	3.70	0.70	0	4.00	0
Rainfall (in):	2.01	0.03	1.12	7.00	1.50	3.38

Trials conducted by Larry Thompson.

Tifton, Georgia:
Earlier Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
AM1511 B2RF	1979	45.7	83.8	1.14	29.5	4.8
PHY 499 WRF	1946	46.1	85.3	1.14	34.5	4.4
DP 1028 B2RF	1918	47.5	85.0	1.14	28.8	4.6
CG 3787 B2RF	1914	45.7	85.1	1.15	30.2	4.7
DP 0924 B2RF	1857	44.4	84.8	1.13	31.4	5.0
DP 0920 B2RF	1845	44.9	84.9	1.16	28.8	4.5
GA2004143	1841	44.3	85.6	1.25	34.6	4.1
DP 0912 B2RF	1830	43.1	84.4	1.13	30.5	4.6
FM1740B2RF	1813	44.1	84.6	1.14	29.6	4.2
ST 4145LLB2	1806	43.1	84.8	1.16	34.3	3.9
PHY 375 WRF	1783	44.6	84.3	1.14	30.2	4.3
BCSX 1150B2RF	1781	40.7	85.5	1.20	34.9	4.6
PHY 367 WRF	1778	43.3	84.8	1.16	30.5	4.2
BRS293	1769	41.5	83.8	1.15	34.5	4.6
Dyna-Gro 2570B2RF	1765	42.3	84.4	1.15	30.9	4.3
DP 0949B2RF	1752	45.1	84.6	1.17	32.5	4.9
DP 1133 B2RF	1752	46.0	85.4	1.16	33.3	4.4
All-Tex ATX81144	1751	42.1	85.3	1.22	32.8	4.0
All-Tex 7A21	1745	45.0	84.8	1.16	31.8	4.9
ST 4288B2F	1737	42.0	83.7	1.14	27.3	4.9
BX 1262B2F	1732	43.3	84.5	1.19	31.4	4.3
AM 1550 B2RF	1718	42.1	84.6	1.14	29.1	4.1
All-Tex ATX3039 B2RF	1713	44.5	83.8	1.16	27.7	4.5
GA2006106	1677	42.7	84.4	1.18	32.4	4.4
SSG CT Linwood	1664	43.4	85.2	1.10	33.7	4.9
BRS286	1634	40.7	82.8	1.10	31.9	4.5
SSG HQ 210 CT	1629	39.9	83.9	1.16	32.7	4.7
All-Tex LA122	1622	43.9	84.1	1.15	28.8	4.4
BX 1252LLB2	1605	44.1	84.3	1.18	31.1	4.5
SSG CT310 HQ	1526	40.6	84.3	1.13	34.2	4.5
GA2008057	1501	41.5	85.0	1.22	33.8	4.1
Average	1754	43.5	84.6	1.16	31.5	4.5
LSD 0.10	137	0.9	1.1	0.04	2.3	0.4
CV %	6.6	1.7	0.8	1.97	4.3	5.4

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: April 27, 2011.

Harvested: September 12, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 78 lb N, 54 lb P₂O₅, and 108 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0.50	2.30	3.00	1.00	1.00	0
Rainfall (in):	1.48	0	1.94	4.06	1.26	4.53

Trials conducted by Larry Thompson.

Tifton, Georgia:
Earlier Maturity Cotton Variety Performance
including Micro-Gin Quality Data, 2011, Irrigated

Variety	Lint Yield	MG ¹ Lint Yield	Lint	MG ¹ Lint	Unif. Index ²	MG ¹ Unif. Index	Length ²	MG ¹ Length	Strength ²	MG ¹ Strength*	Mic. ²	MG ¹ Mic.
	lb/acre	lb/acre	%	%	%	%	inches	inches	g/tex	g/tex	units	units
All-Tex 7A21	1745	1588	45.0	40.0	84.8	83.4	1.16	1.17	31.8	31.7	4.9	5.0
All-Tex ATX3039 B2RF	1713	1549	44.5	39.6	83.8	82.5	1.16	1.15	27.7	28.2	4.5	4.5
All-Tex ATX81144	1751	1576	42.1	37.7	85.3	84.3	1.22	1.23	32.8	33.4	4.0	4.1
All-Tex LA122	1622	1522	43.9	40.5	84.1	83.0	1.15	1.16	28.8	27.9	4.4	4.5
AM 1550 B2RF	1718	1646	42.1	39.9	84.6	82.0	1.14	1.13	29.1	28.5	4.1	4.8
AM1511 B2RF	1979	1630	45.7	41.1	83.8	82.7	1.14	1.13	29.5	30.1	4.8	5.1
BCSX 1150B2RF	1781	1582	40.7	36.2	85.5	84.4	1.20	1.21	34.9	33.3	4.6	4.7
BRS286	1634	1524	40.7	38.0	82.8	82.0	1.10	1.12	31.9	30.7	4.5	4.7
BRS293	1769	1618	41.5	37.6	83.8	82.9	1.15	1.16	34.5	33.7	4.6	4.8
BX 1252LLB2	1605	1433	44.1	38.9	84.3	82.9	1.18	1.15	31.1	32.6	4.5	5.0
BX 1262B2F	1732	1451	43.3	39.2	84.5	83.2	1.19	1.16	31.4	30.6	4.3	4.8
CG 3787 B2RF	1914	1625	45.7	42.3	85.1	83.4	1.15	1.16	30.2	29.5	4.7	4.9
DP 0912 B2RF	1830	1683	43.1	39.4	84.4	82.4	1.13	1.11	30.5	29.3	4.6	5.3
DP 0920 B2RF	1845	1703	44.9	41.1	84.9	82.5	1.16	1.16	28.8	28.3	4.5	5.1
DP 0924 B2RF	1857	1647	44.4	38.8	84.8	82.8	1.13	1.12	31.4	30.3	5.0	5.2
DP 0949B2RF	1752	1616	45.1	40.9	84.6	82.7	1.17	1.14	32.5	30.5	4.9	5.2
DP 1028 B2RF	1918	1771	47.5	43.3	85.0	83.5	1.14	1.13	28.8	29.1	4.6	5.0
DP 1133 B2RF	1752	1609	46.0	41.8	85.4	84.3	1.16	1.16	33.3	31.9	4.4	5.0
Dyna-Gro 2570B2RF	1765	1700	42.3	39.9	84.4	83.0	1.15	1.15	30.9	29.9	4.3	4.8
FM1740B2RF	1813	1534	44.1	39.7	84.6	83.1	1.14	1.14	29.6	29.7	4.2	4.8
GA2004143	1841	1712	44.3	40.9	85.6	83.3	1.25	1.21	34.6	32.0	4.1	4.6
GA2006106	1677	1528	42.7	38.0	84.4	83.2	1.18	1.19	32.4	32.6	4.4	4.8
GA2008057	1501	1358	41.5	36.8	85.0	83.6	1.22	1.19	33.8	33.3	4.1	4.5
PHY 367 WRF	1778	1632	43.3	39.5	84.8	83.0	1.16	1.16	30.5	29.5	4.2	4.5
PHY 375 WRF	1783	1615	44.6	40.3	84.3	82.5	1.14	1.14	30.2	28.5	4.3	4.6
PHY 499 WRF	1946	1783	46.1	41.9	85.3	83.3	1.14	1.13	34.5	32.9	4.4	4.8
SSG CT Linwood	1664	1517	43.4	39.1	85.2	83.1	1.10	1.12	33.7	32.9	4.9	5.2
SSG CT310 HQ	1526	1429	40.6	36.3	84.3	83.1	1.13	1.15	34.2	34.8	4.5	4.7
SSG HQ 210 CT	1629	1567	39.9	37.8	83.9	81.8	1.16	1.13	32.7	31.5	4.7	5.0
ST 4145LLB2	1806	1596	43.1	37.7	84.8	83.4	1.16	1.17	34.3	31.1	3.9	4.6
ST 4288B2F	1737	1532	42.0	36.9	83.7	82.3	1.14	1.16	27.3	28.7	4.9	4.9
Average	1754	1590	43.5	39.4	84.6	83.0	1.16	1.15	31.5	30.9	4.5	4.8
LSD 0.10	137	175	0.9	0.7	1.1	0.6	0.04	0.02	2.3	1.2	0.4	0.2
CV %	6.6	9.3	1.7	1.5	0.8	0.6	1.97	1.71	4.3	3.3	5.4	2.9

1. Micro-Gin quality samples are from total seed cotton harvested from each plot.
2. A random quality sample was taken on the picker during cotton harvest.
3. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: April 27, 2011.

Harvested: September 12, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 78 lb N, 54 lb P₂O₅, and 108 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0.50	2.30	3.00	1.00	1.00	0
Rainfall (in):	1.48	0	1.94	4.06	1.26	4.53

Trials conducted by Larry Thompson.

Yield Summary for Earlier Maturity Cotton Varieties, 2011, Irrigated

Variety	Lint Yield ^a					Lint %	Unif. Index %	Length in	Strength g/tex	Mic. units
	Bainbridge	Midville	Plains lb/acre	Tifton	4-Loc. Average					
DP 0912 B2RF	1847 ⁵	2924 ²	2022 ²	1830 ⁸	2156 ¹	43.7	84.0	1.14	29.5	4.7
PHY 499 WRF	1877 ⁴	2957 ¹	1778 ¹⁴	1946 ²	2139 ²	46.5	84.9	1.15	32.4	4.6
AM1511 B2RF	1886 ³	2637 ⁵	1957 ³	1979 ¹	2115 ³	45.7	84.4	1.15	29.9	4.7
DP 1028 B2RF	1942 ¹	2777 ³	1664 ²⁰	1918 ³	2075 ⁴	46.2	85.1	1.15	28.6	4.8
CG 3787 B2RF	1781 ⁶	2547 ¹²	1908 ⁸	1914 ⁴	2038 ⁵	44.7	84.9	1.16	29.3	4.6
ST 4145LLB2	1566 ²⁰	2494 ¹⁸	2150 ¹	1806 ¹⁰	2004 ⁶	42.7	84.7	1.18	32.6	4.2
DP 1133 B2RF	1928 ²	2335 ²⁶	1947 ⁴	1752 ^{16T}	1991 ⁷	45.9	85.3	1.17	31.9	4.6
All-Tex ATX3039 B2RF	1593 ¹⁸	2741 ⁴	1912 ⁷	1713 ²²	1990 ⁸	44.1	84.1	1.16	28.5	4.3
PHY 375 WRF	1684 ¹⁰	2528 ¹⁶	1806 ¹²	1783 ¹¹	1950 ⁹	44.2	84.3	1.16	29.0	4.4
BX 1262B2F	1667 ¹²	2534 ^{13T}	1821 ¹¹	1732 ²⁰	1938 ¹⁰	43.3	84.7	1.18	30.8	4.6
BX 1252LLB2	1681 ¹¹	2567 ¹¹	1849 ⁹	1605 ²⁸	1926 ¹¹	42.7	84.3	1.18	30.9	4.5
PHY 367 WRF	1640 ¹⁴	2328 ²⁷	1931 ⁵	1778 ¹³	1919 ¹²	43.2	84.7	1.18	29.7	4.4
Dyna-Gro 2570B2RF	1761 ⁷	2426 ²²	1722 ¹⁷	1765 ¹⁵	1918 ¹³	42.5	84.7	1.17	30.4	4.5
BCSX 1150B2RF	1626 ¹⁵	2580 ¹⁰	1677 ¹⁸	1781 ¹²	1916 ¹⁴	40.1	85.2	1.22	33.7	4.7
DP 0920 B2RF	1507 ²⁵	2510 ¹⁷	1798 ¹³	1845 ⁶	1915 ¹⁵	44.0	84.4	1.16	28.2	4.6
DP 0949B2RF	1655 ¹³	2610 ⁶	1624 ²²	1752 ^{16T}	1910 ¹⁶	44.2	84.7	1.17	30.8	4.8
GA2004143	1617 ¹⁶	2597 ⁸	1577 ²⁴	1841 ⁷	1908 ¹⁷	44.2	85.3	1.23	33.8	4.4
ST 4288B2F	1606 ¹⁷	2341 ²⁵	1924 ⁶	1737 ¹⁹	1902 ¹⁸	40.9	83.9	1.17	28.1	4.7
FM1740B2RF	1580 ¹⁹	2532 ¹⁴	1667 ¹⁹	1813 ⁹	1898 ¹⁹	43.3	84.0	1.16	29.1	4.4
All-Tex 7A21	1503 ²⁶	2490 ¹⁹	1848 ¹⁰	1745 ¹⁸	1897 ²⁰	44.2	84.8	1.19	31.1	4.6
DP 0924 B2RF	1468 ²⁸	2599 ⁷	1661 ²¹	1857 ⁵	1896 ²¹	43.1	84.7	1.15	30.6	4.8
AM 1550 B2RF	1526 ²²	2439 ²¹	1752 ¹⁶	1718 ²¹	1859 ²²	42.1	84.3	1.15	28.3	4.5
All-Tex ATX81144	1481 ²⁷	2530 ¹⁵	1595 ²³	1751 ¹⁷	1839 ²³	42.0	85.1	1.24	32.2	3.9
All-Tex LA122	1524 ²³	2387 ²³	1772 ¹⁵	1622 ²⁷	1826 ²⁴	44.2	84.8	1.17	29.1	4.4
GA2006106	1691 ⁹	2534 ^{13T}	1358 ²⁸	1677 ²³	1815 ²⁵	42.1	84.9	1.22	33.3	4.4
BRS293	1542 ²¹	2582 ⁹	1246 ³⁰	1769 ¹⁴	1785 ²⁶	41.7	84.1	1.17	34.2	4.6
SSG HQ 210 CT	1399 ²⁹	2451 ²⁰	1541 ²⁶	1629 ²⁶	1755 ²⁷	40.6	83.4	1.15	31.6	4.6
BRS286	1707 ⁸	2070 ²⁹	1567 ²⁵	1634 ²⁵	1745 ²⁸	41.5	83.4	1.13	31.7	4.5
SSG CT Linwood	1185 ³¹	2347 ²⁴	1256 ²⁹	1664 ²⁴	1613 ²⁹	42.6	84.8	1.12	33.6	5.0
SSG CT310 HQ	1522 ²⁴	2197 ²⁸	1194 ³¹	1526 ²⁹	1610 ³⁰	40.6	84.3	1.15	33.9	4.7
GA2008057	1302 ³⁰	1750 ³⁰	1452 ²⁷	1501 ³⁰	1501 ³¹	41.1	85.1	1.21	33.6	4.2
Average	1622	2495	1709	1754	1895	43.2	84.6	1.17	31.0	4.5
LSD 0.10	190	244	270	137	170	0.9	0.6	0.02	1.1	0.2
CV %	10.0	8.4	13.4	6.6	9.7	2.1	0.8	1.95	4.7	5.0

^a Superscripts indicate ranking at that location.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

**Two-Year Summary for Earlier Maturity Cotton Varieties
at Four Locations^a, 2010-2011, Irrigated**

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length inches	Strength g/tex	Micronaire units
			Index %			
PHY 499 WRF	1982	46.1	84.9	1.15	32.5	4.7
DP 0912 B2RF	1958	43.1	83.8	1.12	30.0	4.8
DP 1028 B2RF	1885	46.1	84.7	1.16	29.4	4.8
DP 0924 B2RF	1785	43.1	83.9	1.13	30.5	4.9
FM1740B2RF	1776	43.3	83.8	1.15	29.7	4.6
Dyna-Gro 2570B2RF	1773	42.5	84.4	1.16	30.3	4.6
PHY 375 WRF	1771	44.1	84.1	1.16	29.7	4.4
DP 0920 B2RF	1762	43.8	84.2	1.15	28.4	4.6
PHY 367 WRF	1752	42.9	84.1	1.17	30.4	4.2
AM 1550 B2RF	1740	42.6	83.8	1.13	28.4	4.5
ST 4288B2F	1726	40.6	83.5	1.16	28.5	4.7
All-Tex 7A21	1712	44.0	84.7	1.19	31.5	4.5
All-Tex LA122	1675	44.0	84.7	1.18	29.4	4.4
GA2006106	1603	41.7	84.6	1.22	33.5	4.4
SSG CT Linwood	1553	42.8	84.3	1.12	33.4	5.0
Average	1764	43.4	84.2	1.16	30.4	4.6
LSD 0.10	78	0.4	0.4	0.01	0.7	0.1
CV %	10.6	2.0	0.7	1.85	4.2	5.1

^a Bainbridge, Midville, Plains, and Tifton.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Bainbridge, Georgia: Later Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield	Lint	Uniformity Index*	Length*	Strength*	Micronaire*
	lb/acre	%	%	inches	g/tex	units
PHY 499 WRF	1932	45.3	84.5	1.15	32.5	4.7
DP 1252 B2RF	1782	46.6	85.4	1.16	30.0	5.1
ST 4145LLB2	1774	42.4	85.2	1.16	31.8	4.9
BX 1262B2F	1771	42.5	84.4	1.17	30.8	5.0
MON 10R051 B2RF	1749	44.9	85.6	1.18	29.2	4.9
DP 1137 B2RF	1727	45.5	84.8	1.15	28.3	5.1
BX 1252LLB2	1708	44.0	85.2	1.17	31.7	5.1
ST 5288B2F	1708	43.1	84.3	1.13	27.8	5.1
DP 1050 B2RF	1693	45.3	84.6	1.15	28.8	4.9
AM1511 B2RF	1665	43.7	84.2	1.14	29.3	4.7
GA2007095	1663	41.2	85.3	1.19	33.0	5.1
DP 1133 B2RF	1636	44.7	85.3	1.17	30.9	4.9
GA2004230	1636	42.0	84.8	1.22	31.9	4.8
GA2008083	1631	44.8	83.4	1.15	30.2	5.0
DP 1048 B2RF	1584	43.1	85.0	1.17	29.2	4.7
DP 1034 B2RF	1537	46.3	84.9	1.18	28.6	5.0
PHY 565 WRF	1519	42.9	84.4	1.15	30.1	4.8
PHY 375 WRF	1505	44.8	84.4	1.16	29.2	4.9
ST 5458B2RF	1484	43.8	84.5	1.16	30.3	5.2
BX 1261B2F	1433	41.4	84.9	1.18	29.6	4.5
BX 1254LLB2	1394	43.6	84.4	1.19	31.2	5.2
PHY 440 W	1193	41.2	84.8	1.15	31.2	4.8
Average	1624	43.8	84.7	1.16	30.2	4.9
LSD 0.10	157	1.0	0.9	N.S. ¹	2.0	0.3
CV %	8.2	2.0	0.6	1.87	3.8	3.2

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 11, 2011.

Harvested: October 17, 2011.

Soil Type: Bonnaeu blanton loamy sand.

Fertilization: 180 lb N, 71 lb P₂O₅, and 122 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	May	June	July	Aug.	Sept.
Irrigation (in):	1.60	4.80	1.60	3.20	0.80
Rainfall (in):	2.10	1.35	9.10	2.35	5.05

Trials conducted by Larry Thompson.

Midville, Georgia:
Later Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield	Lint	Uniformity Index*	Length*	Strength*	Micronaire*
	lb/acre	%	%	inches	g/tex	units
PHY 499 WRF	3021	45.7	85.2	1.16	32.5	4.6
DP 1252 B2RF	2864	48.3	84.8	1.17	27.7	4.7
DP 1137 B2RF	2863	46.3	84.7	1.15	26.7	4.7
MON 10R051 B2RF	2794	47.4	85.2	1.19	25.6	4.8
AM1511 B2RF	2786	46.5	84.6	1.16	30.7	4.9
DP 1050 B2RF	2769	46.8	85.0	1.19	28.6	4.7
BX 1252LLB2	2754	44.3	83.9	1.18	30.9	4.7
ST 5458B2RF	2710	42.8	84.2	1.17	29.9	5.0
PHY 375 WRF	2674	44.9	83.5	1.15	29.1	4.1
BX 1261B2F	2652	42.5	84.4	1.17	28.8	4.6
GA2008083	2646	46.9	83.1	1.14	29.1	4.6
BX 1262B2F	2638	43.6	84.9	1.18	31.2	4.7
DP 1034 B2RF	2624	46.3	85.1	1.20	26.5	4.6
GA2007095	2576	42.7	84.0	1.19	29.5	4.9
DP 1048 B2RF	2570	46.1	85.0	1.17	27.4	4.8
GA2004230	2524	42.5	84.8	1.24	29.8	4.5
PHY 565 WRF	2509	43.0	85.1	1.16	32.6	4.7
BX 1254LLB2	2503	44.9	83.6	1.16	29.7	5.3
ST 4145LLB2	2456	42.2	84.5	1.16	30.4	4.5
ST 5288B2F	2425	43.0	83.2	1.15	27.4	4.5
DP 1133 B2RF	2410	45.3	85.6	1.20	30.9	4.7
PHY 440 W	2383	42.4	84.6	1.15	29.0	4.4
Average	2643	44.7	84.5	1.17	29.2	4.7
LSD 0.10	227	0.9	1.3	0.04	1.7	0.4
CV %	7.3	1.6	0.9	1.76	3.7	4.4

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 3, 2011.

Harvested: October 3, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 96 lb N, 80 lb P₂O₅, and 60 lb K₂O/acre.

Management: Temik applied 5 lb/acre and Telone II applied 3 gal/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0	1.90	2.45	4.00	4.00	0
Rainfall (in):	1.41	2.32	1.76	2.22	1.50	2.51

Trials conducted by Larry Thompson.

Plains, Georgia:
Later Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
AM1511 B2RF	1995	43.6	85.3	1.17	31.1	4.6
DP 1048 B2RF	1957	42.3	84.9	1.21	28.9	4.0
DP 1050 B2RF	1840	45.7	85.8	1.21	29.0	4.2
MON 10R051 B2RF	1838	44.4	85.5	1.20	29.0	4.2
DP 1252 B2RF	1778	44.2	85.0	1.19	30.4	4.1
DP 1034 B2RF	1777	43.8	86.1	1.21	28.9	4.4
DP 1133 B2RF	1768	44.6	86.0	1.20	32.7	4.4
ST 4145LLB2	1755	40.5	84.8	1.19	32.8	4.1
PHY 499 WRF	1689	42.8	85.7	1.19	32.4	4.2
DP 1137 B2RF	1682	43.3	85.4	1.18	29.8	4.1
ST 5288B2F	1637	39.5	84.7	1.20	30.1	4.3
BX 1262B2F	1618	41.1	85.6	1.21	31.4	4.1
BX 1254LLB2	1584	42.2	85.1	1.21	32.5	4.5
GA2004230	1571	40.2	85.6	1.28	33.1	4.2
ST 5458B2RF	1562	39.3	84.9	1.22	33.1	4.3
GA2007095	1545	41.6	84.5	1.19	31.8	4.1
PHY 375 WRF	1491	41.6	84.9	1.20	30.6	3.8
BX 1252LLB2	1486	37.6	85.6	1.22	31.7	4.3
PHY 440 W	1480	40.2	85.0	1.20	30.6	4.3
PHY 565 WRF	1368	40.6	85.8	1.21	32.9	3.9
BX 1261B2F	1351	39.3	84.7	1.22	30.5	3.8
GA2008083	1341	41.8	86.4	1.24	34.3	4.4
Average	1642	41.8	85.3	1.21	31.2	4.2
LSD 0.10	213	1.5	0.9	0.03	1.6	0.4
CV %	11.0	3.0	0.6	1.68	2.9	5.1

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 5, 2011.

Harvested: October 7, 2011.

Soil Type: Greenville sandy clay loam.

Fertilization: 100 lb N, 88 lb P₂O₅, and 24 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0	3.70	0.70	0	4.00	0
Rainfall (in):	2.01	0.03	1.12	7.00	1.50	3.38

Trials conducted by Larry Thompson.

Tifton, Georgia:
Later Maturity Cotton Variety Performance, 2011, Irrigated

Variety	Lint Yield	Lint	Uniformity Index*	Length*	Strength*	Micronaire*
	lb/acre	%	%	inches	g/tex	units
PHY 499 WRF	2072	44.8	84.6	1.13	32.4	4.3
DP 1050 B2RF	2027	46.4	85.2	1.19	27.2	4.6
AM1511 B2RF	1976	45.8	84.8	1.15	29.7	4.6
DP 1048 B2RF	1965	45.6	85.0	1.18	28.9	4.4
DP 1252 B2RF	1958	45.5	85.2	1.18	28.6	4.2
DP 1034 B2RF	1933	46.9	84.6	1.17	26.5	4.6
PHY 375 WRF	1901	43.7	84.6	1.17	28.8	4.1
ST 4145LLB2	1897	41.6	85.9	1.19	32.8	3.8
DP 1137 B2RF	1887	44.8	85.3	1.17	29.4	4.3
MON 10R051 B2RF	1877	46.1	85.8	1.17	29.7	4.5
BX 1262B2F	1853	42.2	84.5	1.18	31.1	4.2
DP 1133 B2RF	1848	45.7	85.4	1.17	32.3	4.4
ST 5288B2F	1782	42.9	84.1	1.14	29.8	4.3
ST 5458B2RF	1772	41.6	83.6	1.17	32.5	4.6
BX 1254LLB2	1770	42.6	84.8	1.19	33.2	4.6
PHY 565 WRF	1767	41.4	85.0	1.21	32.0	3.9
BX 1252LLB2	1732	41.7	85.2	1.20	31.6	4.0
GA2007095	1729	42.4	84.2	1.17	30.9	4.3
GA2004230	1709	43.1	83.5	1.22	30.9	4.3
BX 1261B2F	1677	39.1	85.2	1.16	30.9	3.9
GA2008083	1641	43.9	84.9	1.19	31.7	4.3
PHY 440 W	1591	40.7	84.6	1.15	32.2	4.3
Average	1835	43.6	84.8	1.17	30.6	4.3
LSD 0.10	138	1.2	1.1	0.04	2.3	N.S. ¹
CV %	6.4	2.3	0.8	2.11	4.4	6.8

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: April 27, 2011.

Harvested: September 12, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 78 lb N, 54 lb P₂O₅, and 108 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0.50	2.30	3.00	1.00	1.00	0
Rainfall (in):	1.48	0	1.94	4.06	1.26	4.53

Trials conducted by Larry Thompson.

Tifton, Georgia:
Later Maturity Cotton Variety Performance
including Micro-Gin Quality Data, 2011, Irrigated

Variety	Lint Yield lb/acre	MG ¹ Lint Yield lb/acre	Lint %	MG ¹ Lint %	Unif. Index ² %	MG ¹ Unif. Index %	Length ² inches	MG ¹ Length inches	Strength ² g/tex	MG ¹ Strength* g/tex	Mic. ² units	MG ¹ Mic. units
AM1511 B2RF	1976	1768	45.8	40.7	84.8	83.2	1.15	1.15	29.7	31.4	4.6	4.9
BX 1252LLB2	1732	1633	41.7	38.2	85.2	83.3	1.20	1.18	31.6	32.4	4.0	4.7
BX 1254LLB2	1770	1670	42.6	39.1	84.8	83.1	1.19	1.18	33.2	32.5	4.6	5.0
BX 1261B2F	1677	1629	39.1	36.9	85.2	83.3	1.16	1.16	30.9	31.9	3.9	4.4
BX 1262B2F	1853	1766	42.2	39.0	84.5	83.3	1.18	1.17	31.1	31.8	4.2	4.7
DP 1034 B2RF	1933	1760	46.9	41.6	84.6	83.4	1.17	1.17	26.5	29.3	4.6	4.7
DP 1048 B2RF	1965	1865	45.6	41.8	85.0	83.4	1.18	1.17	28.9	28.7	4.4	4.7
DP 1050 B2RF	2027	1923	46.4	42.7	85.2	83.2	1.19	1.17	27.2	28.1	4.6	4.6
DP 1133 B2RF	1848	1717	45.7	41.3	85.4	84.1	1.17	1.16	32.3	32.6	4.4	4.8
DP 1137 B2RF	1887	1817	44.8	41.9	85.3	82.5	1.17	1.13	29.4	29.1	4.3	4.9
DP 1252 B2RF	1958	1922	45.5	43.4	85.2	83.0	1.18	1.14	28.6	29.6	4.2	4.8
GA2004230	1709	1576	43.1	38.6	83.5	83.9	1.22	1.24	30.9	31.8	4.3	4.4
GA2007095	1729	1611	42.4	38.2	84.2	83.3	1.17	1.18	30.9	31.8	4.3	4.4
GA2008083	1641	1526	43.9	39.6	84.9	83.0	1.19	1.18	31.7	32.4	4.3	4.7
MON 10R051 B2RF	1877	1762	46.1	42.1	85.8	83.3	1.17	1.15	29.7	29.0	4.5	4.7
PHY 375 WRF	1901	1769	43.7	39.4	84.6	82.7	1.17	1.15	28.8	29.3	4.1	4.6
PHY 440 W	1591	1501	40.7	37.2	84.6	83.3	1.15	1.15	32.2	31.3	4.3	4.5
PHY 499 WRF	2072	1966	44.8	41.5	84.6	83.5	1.13	1.15	32.4	31.6	4.3	4.9
PHY 565 WRF	1767	1654	41.4	37.7	85.0	83.4	1.21	1.19	32.0	32.0	3.9	4.5
ST 4145LLB2	1897	1730	41.6	37.1	85.9	83.4	1.19	1.17	32.8	31.8	3.8	4.4
ST 5288B2F	1782	1642	42.9	38.3	84.1	82.6	1.14	1.16	29.8	28.6	4.3	4.8
ST 5458B2RF	1772	1682	41.6	38.2	83.6	83.2	1.17	1.18	32.5	32.1	4.6	5.1
Average	1835	1722	43.6	39.8	84.8	83.2	1.17	1.17	30.6	30.9	4.3	4.7
LSD 0.10	138	117	1.2	0.5	1.1	0.5	0.04	0.02	2.3	1.1	N.S. ¹	0.2
CV %	6.4	5.8	2.3	1.1	0.8	0.5	2.11	1.43	4.4	2.9	6.8	3.6

1. Micro-Gin quality samples are from total seed cotton harvested from each plot.

2. A random quality sample was taken on the picker during cotton harvest.

3. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: April 27, 2011.

Harvested: September 12, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 78 lb N, 54 lb P₂O₅, and 108 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

April May June July Aug. Sept.

Irrigation (in): 0.50 2.30 3.00 1.00 1.00 0

Rainfall (in): 1.48 0 1.94 4.06 1.26 4.53

Trials conducted by Larry Thompson.

Yield Summary for Later Maturity Cotton Varieties, 2011, Irrigated

Variety	Lint Yield ^a					Lint %	Unif. Index %	Length in	Strength g/tex	Mic. units
	Bainbridge	Midville	Plains lb/acre	Tifton	4-Loc. Average					
PHY 499 WRF	1932 ¹	3021 ¹	1689 ⁹	2072 ¹	2179 ¹	44.7	85.0	1.15	32.4	4.4
AM1511 B2RF	1665 ⁹	2786 ⁵	1995 ¹	1976 ³	2105 ²	44.9	84.7	1.15	30.2	4.7
DP 1252 B2RF	1782 ²	2864 ²	1778 ⁵	1958 ⁵	2095 ³	46.2	85.1	1.17	29.2	4.5
DP 1050 B2RF	1693 ⁸	2769 ⁶	1840 ³	2027 ²	2082 ⁴	46.0	85.1	1.18	28.4	4.6
MON 10R051 B2RF	1749 ⁵	2794 ⁴	1838 ⁴	1877 ¹⁰	2064 ⁵	45.7	85.5	1.18	28.3	4.6
DP 1137 B2RF	1727 ⁶	2863 ³	1682 ¹⁰	1887 ⁹	2040 ⁶	45.0	85.0	1.16	28.5	4.5
DP 1048 B2RF	1584 ¹³	2570 ¹⁵	1957 ²	1965 ⁴	2019 ⁷	44.3	85.0	1.18	28.6	4.4
ST 4145LLB2	1774 ³	2456 ¹⁹	1755 ⁸	1897 ⁸	1971 ⁸	41.7	85.1	1.17	31.9	4.3
BX 1262B2F	1771 ⁴	2638 ¹²	1618 ¹²	1853 ¹¹	1970 ⁹	42.3	84.8	1.18	31.1	4.5
DP 1034 B2RF	1537 ¹⁴	2624 ¹³	1777 ⁶	1933 ⁶	1968 ¹⁰	45.8	85.1	1.19	27.6	4.6
BX 1252LLB2	1708 ^{7T}	2754 ⁷	1486 ¹⁸	1732 ¹⁷	1920 ¹¹	41.9	85.0	1.19	31.5	4.5
DP 1133 B2RF	1636 ^{11T}	2410 ²¹	1768 ⁷	1848 ¹²	1916 ¹²	45.1	85.6	1.18	31.7	4.6
PHY 375 WRF	1505 ¹⁶	2674 ⁹	1491 ¹⁷	1901 ⁷	1893 ¹³	43.8	84.3	1.17	29.4	4.2
ST 5288B2F	1708 ^{7T}	2425 ²⁰	1637 ¹¹	1782 ¹³	1888 ¹⁴	42.1	84.1	1.16	28.8	4.5
ST 5458B2RF	1484 ¹⁷	2710 ⁸	1562 ¹⁵	1772 ¹⁴	1882 ¹⁵	41.9	84.3	1.18	31.4	4.8
GA2007095	1663 ¹⁰	2576 ¹⁴	1545 ¹⁶	1729 ¹⁸	1878 ¹⁶	42.0	84.5	1.18	31.3	4.6
GA2004230	1636 ^{11T}	2524 ¹⁶	1571 ¹⁴	1709 ¹⁹	1860 ¹⁷	41.9	84.7	1.24	31.4	4.4
GA2008083	1631 ¹²	2646 ¹¹	1341 ²²	1641 ²¹	1815 ¹⁸	44.3	84.4	1.18	31.3	4.6
BX 1254LLB2	1394 ¹⁹	2503 ¹⁸	1584 ¹³	1770 ¹⁵	1813 ¹⁹	43.3	84.4	1.19	31.6	4.9
PHY 565 WRF	1519 ¹⁵	2509 ¹⁷	1368 ²⁰	1767 ¹⁶	1791 ²⁰	42.0	85.1	1.18	31.9	4.3
BX 1261B2F	1433 ¹⁸	2652 ¹⁰	1351 ²¹	1677 ²⁰	1778 ²¹	40.6	84.8	1.18	29.9	4.2
PHY 440 W	1193 ²⁰	2383 ²²	1480 ¹⁹	1591 ²²	1662 ²²	41.1	84.8	1.16	30.7	4.4
Average	1624	2643	1642	1835	1936	43.5	84.8	1.18	30.3	4.5
LSD 0.10	157	227	213	138	135	1.1	0.6	0.02	1.1	0.2
CV %	8.2	7.3	11.0	6.4	8.2	2.2	0.7	1.86	3.7	4.9

^a Superscripts indicate ranking at that location.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

**Two-Year Summary for Later Maturity Cotton Varieties
at Four Locations^a, 2010-2011, Irrigated**

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length inches	Strength g/tex	Micronaire units
			Index %			
PHY 499 WRF	1893	45.0	84.6	1.14	32.5	4.6
DP 1252 B2RF	1841	46.1	84.7	1.17	29.3	4.7
DP 1050 B2RF	1840	46.0	84.8	1.17	28.6	4.7
DP 1137 B2RF	1803	45.3	84.5	1.15	29.0	4.7
DP 1034 B2RF	1794	45.9	84.7	1.18	28.0	4.7
DP 1048 B2RF	1794	44.7	84.7	1.17	28.5	4.6
PHY 375 WRF	1744	43.8	84.1	1.16	29.4	4.3
DP 1133 B2RF	1731	45.1	85.3	1.18	32.0	4.7
ST 5458B2RF	1690	42.3	83.9	1.17	31.4	4.9
PHY 565 WRF	1687	42.7	84.5	1.18	32.2	4.2
ST 5288B2F	1684	42.7	83.4	1.15	28.9	4.6
PHY 440 W	1486	41.6	84.1	1.15	30.6	4.5
Average	1749	44.3	84.4	1.16	30.0	4.6
LSD 0.10	67	0.4	0.4	0.01	0.7	0.1
CV %	9.3	2.4	0.7	1.92	3.8	5.2

^a Bainbridge, Midville, Plains, and Tifton.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Midville, Georgia: Cotton Strains Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire*
			Index* %			
DP 1050 B2RF	2973	48.2	85.8	1.18	27.4	4.8
DP 1044 B2RF	2782	42.5	84.3	1.16	29.6	4.6
All-Tex 9C253 B2RF	2750	44.5	85.3	1.17	33.2	5.1
CT11622	2708	44.7	85.4	1.19	30.6	4.2
MON 10R011 B2RF	2687	42.1	85.5	1.21	35.6	4.5
GA2009100	2677	45.0	85.3	1.21	34.6	3.9
MON 11R159 B2RF	2672	45.4	85.0	1.21	34.0	4.9
GA2009037	2586	43.8	83.6	1.20	30.9	4.5
MON 10R020 B2RF	2524	44.1	84.4	1.16	28.6	4.7
MON 10R013 B2RF	2478	43.5	85.6	1.23	31.9	5.0
AMX003 B2RF	2422	46.0	84.7	1.17	29.8	4.8
CT11212	2417	45.4	84.8	1.15	29.2	4.8
GA2009148	2393	42.9	84.7	1.18	34.3	4.9
GA2008016	2370	43.0	84.6	1.16	36.1	4.8
GA2009180	2362	42.3	85.7	1.25	36.2	4.5
PHY 440 W	2282	42.5	84.1	1.17	32.2	4.6
All-Tex 9W2863 B2RF	2274	42.4	85.5	1.25	32.3	4.6
GA2009147	2126	41.1	83.6	1.22	33.1	4.0
Average	2527	43.9	84.9	1.19	32.2	4.6
LSD 0.10	227	1.5	1.2	0.04	1.7	0.5
CV %	7.6	2.9	0.8	2.12	3.0	6.4

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 3, 2011.

Harvested: October 3, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 96 lb N, 80 lb P₂O₅, and 60 lb K₂O/acre.

Management: Temik applied 5 lb/acre and Telone II applied 3 gal/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0	1.90	2.45	4.00	4.00	0
Rainfall (in):	1.41	2.33	1.73	3.15	1.02	2.40

Trials conducted by Larry Thompson.

Plains, Georgia:
Cotton Strains Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
DP 1050 B2RF	1858	43.9	85.2	1.21	28.3	4.3
CT11622	1840	43.2	85.7	1.19	28.4	4.3
MON 10R020 B2RF	1815	41.8	84.1	1.14	28.6	4.7
All-Tex 9C253 B2RF	1697	43.5	85.0	1.18	32.4	4.7
AMX003 B2RF	1658	44.9	85.4	1.21	28.9	4.6
PHY 440 W	1611	41.0	85.7	1.19	29.5	4.1
GA2009100	1571	43.0	85.9	1.21	35.6	4.1
MON 10R011 B2RF	1558	41.6	84.8	1.22	33.4	3.9
CT11212	1550	42.4	85.0	1.17	28.0	4.7
DP 1044 B2RF	1515	39.0	84.2	1.19	28.7	4.0
MON 10R013 B2RF	1477	40.3	84.9	1.24	30.8	4.6
MON 11R159 B2RF	1423	41.5	84.7	1.22	33.6	4.1
GA2009148	1359	41.6	84.6	1.18	32.0	4.8
All-Tex 9W2863 B2RF	1348	39.1	86.0	1.22	33.1	4.5
GA2008016	1274	39.9	85.4	1.20	32.7	4.5
GA2009037	1231	39.6	84.7	1.21	31.1	4.2
GA2009147	1136	39.9	83.5	1.19	32.3	4.1
GA2009180	1090	38.0	84.9	1.21	35.3	4.0
Average	1501	41.4	85.0	1.20	31.2	4.3
LSD 0.10	233	1.5	N.S. ¹	0.04	2.0	0.4
CV %	13.1	3.0	0.9	1.81	3.6	5.4

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 5, 2011.

Harvested: October 7, 2011.

Soil Type: Greenville sandy clay loam.

Fertilization: 100 lb N, 88 lb P₂O₅, and 24 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0	3.70	0.70	0	4.00	0
Rainfall (in):	2.01	0.03	1.12	7.00	1.50	3.38

Trials conducted by Larry Thompson.

Tifton, Georgia:
Cotton Strains Performance, 2011, Irrigated

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
CT11622	2092	45.4	85.1	1.17	27.5	4.4
MON 10R011 B2RF	2082	42.9	84.2	1.17	29.8	4.2
DP 1044 B2RF	2049	42.2	84.3	1.16	26.4	4.1
DP 1050 B2RF	2016	46.1	84.6	1.18	27.5	4.4
GA2009100	1982	44.9	84.5	1.18	31.7	4.2
MON 11R159 B2RF	1966	42.9	84.1	1.21	32.5	4.2
All-Tex 9C253 B2RF	1946	44.7	84.5	1.17	29.4	4.9
MON 10R020 B2RF	1939	42.8	84.3	1.14	26.9	4.6
CT11212	1909	43.2	85.1	1.15	27.0	4.6
MON 10R013 B2RF	1846	42.7	84.7	1.20	28.5	4.7
GA2009148	1845	44.0	84.1	1.16	30.5	4.5
AMX003 B2RF	1802	44.4	84.3	1.18	26.9	4.7
PHY 440 W	1731	41.4	84.8	1.15	30.0	4.5
All-Tex 9W2863 B2RF	1693	42.0	85.1	1.19	30.0	4.4
GA2009037	1692	43.5	83.4	1.16	29.0	5.0
GA2009147	1530	41.2	83.7	1.19	31.4	4.1
GA2008016	1508	41.8	84.4	1.18	32.3	4.4
GA2009180	1459	40.4	84.1	1.19	32.1	4.1
Average	1838	43.1	84.4	1.17	29.4	4.4
LSD 0.10	159	1.0	N.S. ¹	N.S.	2.2	0.4
CV %	7.3	2.0	0.7	2.58	4.4	5.8

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: April 27, 2011.

Harvested: September 14, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 78 lb N, 54 lb P₂O₅, and 108 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	0.50	2.30	3.00	1.00	1.00	0
Rainfall (in):	1.48	0	1.94	4.06	1.26	4.53

Trials conducted by Larry Thompson.

Yield Summary for Cotton Strains, 2011, Irrigated

Variety	Lint Yield ^a				Lint %	Unif. Index %	Length inches	Strength g/tex	Mic. units
	Midville	Plains	Tifton	3-Loc. Average					
	----- lb/acre -----								
DP 1050 B2RF	2973 ¹	1858 ¹	2016 ⁴	2282 ¹	46.1	85.2	1.19	27.7	4.5
CT11622	2708 ⁴	1840 ²	2092 ¹	2213 ²	44.4	85.4	1.18	28.8	4.3
All-Tex 9C253 B2RF	2750 ³	1697 ⁴	1946 ⁷	2131 ³	44.2	84.9	1.17	31.7	4.9
DP 1044 B2RF	2782 ²	1515 ¹⁰	2049 ³	2115 ⁴	41.2	84.3	1.17	28.2	4.2
DP 1219 B2RF	2687 ⁵	1558 ⁸	2082 ²	2109 ⁵	42.2	84.8	1.20	32.9	4.2
MON 10R020 B2RF	2524 ⁹	1815 ³	1939 ⁸	2092 ⁶	42.9	84.2	1.14	28.0	4.6
GA2009100	2677 ⁶	1571 ⁷	1982 ⁵	2077 ⁷	44.3	85.2	1.20	33.9	4.0
MON 11R159 B2RF	2672 ⁷	1423 ¹²	1966 ⁶	2021 ⁸	43.3	84.6	1.21	33.3	4.4
AMX003 B2RF	2422 ¹¹	1658 ⁵	1802 ¹²	1961 ⁹	45.1	84.8	1.19	28.5	4.7
CT11212	2417 ¹²	1550 ⁹	1909 ⁹	1959 ¹⁰	43.7	85.0	1.16	28.0	4.7
DP 1212 B2RF	2478 ¹⁰	1477 ¹¹	1846 ¹⁰	1934 ¹¹	42.2	85.1	1.22	30.4	4.8
PHY 440 W	2282 ¹⁶	1611 ⁶	1731 ¹³	1875 ¹²	41.6	84.8	1.17	30.5	4.4
GA2009148	2393 ¹³	1359 ¹³	1845 ¹¹	1866 ¹³	42.9	84.4	1.17	32.2	4.7
GA2009037	2586 ⁸	1231 ¹⁶	1692 ¹⁵	1836 ¹⁴	42.3	83.9	1.19	30.3	4.5
All-Tex 9W2863 B2RF	2274 ¹⁷	1348 ¹⁴	1693 ¹⁴	1772 ¹⁵	41.2	85.5	1.22	31.8	4.5
GA2008016	2370 ¹⁴	1274 ¹⁵	1508 ¹⁷	1717 ¹⁶	41.6	84.8	1.18	33.7	4.6
GA2009180	2362 ¹⁵	1090 ¹⁸	1459 ¹⁸	1637 ¹⁷	40.2	84.9	1.22	34.5	4.2
GA2009147	2126 ¹⁸	1136 ¹⁷	1530 ¹⁶	1597 ¹⁸	40.7	83.6	1.20	32.3	4.0
Average	2527	1501	1838	1955	42.8	84.7	1.19	30.9	4.4
LSD 0.10	227	233	159	159	1.1	0.6	0.02	1.2	0.3
CV %	7.6	13.1	7.3	9.0	2.7	0.8	2.19	3.7	5.9

^a Superscripts indicate ranking at that location.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Athens, Georgia:

Dryland Earlier Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
AM1511 B2RF	1778	47.9	84.2	1.09	29.8	4.7
BX 1262B2F	1773	46.2	84.2	1.12	31.5	4.5
PHY 499 WRF	1725	48.5	83.8	1.07	32.2	4.8
AM 1550 B2RF	1678	45.9	84.0	1.10	25.5	4.7
ST 4288B2F	1663	44.0	82.9	1.11	26.1	4.7
All-Tex 7A21	1607	46.4	83.9	1.11	29.6	4.6
DP 0912 B2RF	1544	45.3	84.5	1.13	31.7	5.1
BRS293	1512	44.6	82.7	1.09	33.5	5.0
DP 0924 B2RF	1505	44.1	83.9	1.11	29.3	5.1
CG 3787 B2RF	1463	48.6	83.7	1.12	27.8	4.9
Dyna-Gro 2570B2RF	1450	44.9	83.5	1.07	29.3	4.9
All-Tex ATX3039 B2RF	1433	47.3	82.8	1.13	27.2	4.6
DP 1028 B2RF	1395	49.8	84.2	1.12	28.4	4.8
BCSX 1150B2RF	1383	42.0	84.2	1.16	30.9	4.4
BX 1252LLB2	1376	46.5	83.2	1.11	29.5	4.8
ST 4145LLB2	1371	45.7	82.2	1.08	30.3	4.4
DP 0949B2RF	1357	47.3	82.8	1.11	28.7	5.1
All-Tex ATX81144	1356	42.6	85.4	1.21	33.7	3.7
All-Tex LA122	1355	46.5	84.0	1.11	28.3	4.5
GA2004143	1352	47.8	83.7	1.14	29.7	4.7
PHY 375 WRF	1341	47.3	83.1	1.06	28.5	4.4
DP 0920 B2RF	1321	46.4	84.0	1.11	26.7	4.8
PHY 367 WRF	1318	46.0	84.0	1.09	29.7	4.4
SSG HQ 210 CT	1313	43.1	83.3	1.09	32.3	5.0
BRS286	1306	42.8	82.1	1.08	31.0	4.4
GA2006106	1253	43.8	83.8	1.15	31.7	4.5
FM1740B2RF	1241	45.2	83.3	1.13	29.0	4.5
DP 1133 B2RF	1240	48.1	83.7	1.12	30.2	5.0
SSG CT310 HQ	1166	42.2	83.5	1.13	34.7	4.9
GA2008057	1004	44.4	84.2	1.15	32.2	4.6
SSG CT Linwood	872	45.8	83.3	1.08	33.5	5.1
Average	1402	45.7	83.6	1.11	30.1	4.7
LSD 0.10	254	0.7	1.0	0.03	2.2	0.4
CV %	15.4	1.3	0.7	1.76	4.2	4.7

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 5, 2011.

Harvested: September 19, 2011.

Soil Type: Cecil coarse sandy loam.

Fertilization: 180 lb N, 80 lb P₂O₅, and 80 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Rainfall (in):	2.50	1.40	3.69	1.90	1.00	1.63

Trials conducted by Larry Thompson.

Midville, Georgia:
Dryland Earlier Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
PHY 499 WRF	1238	48.3	83.4	1.07	29.5	4.6
AM1511 B2RF	1123	47.5	83.3	1.04	30.3	4.7
DP 0912 B2RF	1116	44.9	82.6	1.10	27.9	4.7
ST 4145LLB2	1063	45.8	82.5	1.10	29.6	4.3
DP 1028 B2RF	1039	47.3	82.5	1.11	26.9	4.9
All-Tex ATX81144	1036	43.2	84.6	1.20	33.0	4.0
DP 0924 B2RF	1011	43.2	82.5	1.05	28.3	4.5
BX 1262B2F	1002	43.2	82.3	1.10	31.6	4.4
AM 1550 B2RF	997	46.2	83.5	1.08	26.7	4.6
CG 3787 B2RF	994	47.7	84.2	1.10	28.6	4.9
All-Tex 7A21	989	45.0	83.5	1.12	31.2	4.8
BRS293	986	42.4	83.3	1.08	33.9	5.0
ST 4288B2F	982	42.4	83.4	1.12	28.2	4.4
Dyna-Gro 2570B2RF	961	44.4	83.8	1.10	30.4	4.6
PHY 367 WRF	958	45.7	83.3	1.12	28.8	4.4
DP 0949B2RF	949	46.3	83.3	1.10	30.7	4.9
PHY 375 WRF	949	45.7	83.1	1.06	27.5	4.3
All-Tex ATX3039 B2RF	939	47.1	80.9	1.05	25.6	4.5
All-Tex LA122	916	45.7	82.3	1.08	27.7	4.7
BCSX 1150B2RF	888	42.5	83.6	1.13	31.2	4.3
BX 1252LLB2	886	43.4	82.9	1.08	30.2	4.6
GA2004143	874	45.8	83.2	1.14	31.5	4.6
SSG HQ 210 CT	862	42.2	81.4	1.08	31.1	5.0
BRS286	859	40.9	81.8	1.07	28.7	4.4
DP 0920 B2RF	840	44.2	82.7	1.08	25.1	4.4
FM1740B2RF	816	43.3	83.5	1.09	27.1	4.3
DP 1133 B2RF	793	47.4	83.5	1.10	30.1	5.0
SSG CT Linwood	761	43.2	81.7	1.07	30.0	5.3
SSG CT310 HQ	754	41.7	82.5	1.08	32.1	5.1
GA2006106	754	40.8	83.2	1.14	30.3	4.2
GA2008057	621	40.3	84.3	1.12	34.1	4.5
Average	934	44.4	83.0	1.09	29.6	4.6
LSD 0.10	128	0.9	1.3	0.04	2.1	0.4
CV %	11.7	1.7	0.9	2.50	4.1	4.7

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 3, 2011.

Harvested: October 3, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 85 lb N, 60 lb P₂O₅, and 70 lb K₂O/acre.

Management: Temik applied 5 lb/acre and Telone II applied 3 gal/acre.

	April	May	June	July	Aug.	Sept.
Rainfall (in):	1.41	2.33	1.73	3.15	1.02	2.40

Trials conducted by Larry Thompson.

Plains, Georgia:
Dryland Earlier Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
DP 0912 B2RF	687	40.3	84.3	1.12	30.6	4.3
DP 1028 B2RF	672	45.0	84.4	1.13	28.0	4.1
AM1511 B2RF	648	41.5	83.6	1.10	30.9	4.1
PHY 367 WRF	645	40.2	83.1	1.12	30.5	3.8
BX 1252LLB2	627	37.6	83.1	1.14	31.6	3.7
GA2006106	618	40.9	82.7	1.13	31.8	4.0
BRS286	614	39.1	83.0	1.09	31.1	4.1
DP 0920 B2RF	607	39.3	83.7	1.13	26.0	3.7
AM 1550 B2RF	606	38.8	83.8	1.12	27.6	3.7
PHY 499 WRF	604	41.4	82.7	1.13	31.0	3.9
SSG HQ 210 CT	588	38.7	81.4	1.08	29.8	4.3
CG 3787 B2RF	581	43.2	83.7	1.12	28.5	3.9
All-Tex LA122	578	41.3	83.0	1.11	27.5	3.9
FM1740B2RF	567	40.7	82.4	1.10	28.3	3.8
DP 0949B2RF	566	40.3	83.1	1.10	29.1	4.0
GA2004143	555	43.2	83.4	1.14	33.6	4.1
Dyna-Gro 2570B2RF	553	37.7	83.3	1.12	28.9	3.7
DP 1133 B2RF	542	40.3	84.1	1.13	30.1	3.9
BCSX 1150B2RF	534	37.2	83.0	1.16	32.0	3.6
SSG CT Linwood	531	41.3	83.0	1.07	31.2	4.6
DP 0924 B2RF	528	38.9	84.1	1.09	27.8	4.1
SSG CT310 HQ	519	36.6	82.9	1.11	32.5	3.8
ST 4145LLB2	518	39.4	82.9	1.12	29.1	3.8
All-Tex ATX81144	511	38.3	83.5	1.19	31.4	3.3
ST 4288B2F	500	37.1	83.3	1.12	27.5	3.6
PHY 375 WRF	476	39.7	83.1	1.10	28.1	3.5
All-Tex 7A21	475	40.4	83.5	1.15	30.0	4.1
BRS293	459	38.7	82.2	1.08	32.2	4.4
All-Tex ATX3039 B2RF	431	40.2	82.5	1.11	26.0	3.6
BX 1262B2F	381	37.1	83.0	1.14	31.6	3.4
GA2008057	317	38.0	83.8	1.18	33.3	3.9
Average	550	39.8	83.2	1.12	29.9	3.9
LSD 0.10	N.S. ¹	1.3	1.0	0.03	2.4	0.4
CV %	31.1	2.8	0.7	1.68	4.6	5.5

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 5, 2011.

Harvested: October 7, 2011.

Soil Type: Greenville sandy clay loam.

Fertilization: 100 lb N, 88 lb P₂O₅, and 24 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Rainfall (in):	2.01	0.03	1.12	7.00	1.50	3.38

Trials conducted by Larry Thompson.

Tifton, Georgia:
Dryland Earlier Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
DP 1028 B2RF	2130	46.9	85.2	1.16	28.5	4.4
PHY 499 WRF	2067	46.7	85.7	1.15	32.0	4.4
DP 0920 B2RF	1986	45.2	85.0	1.21	30.0	4.3
DP 0912 B2RF	1964	43.4	84.0	1.13	29.5	4.9
AM1511 B2RF	1946	45.1	84.2	1.13	30.3	4.7
Dyna-Gro 2570B2RF	1916	43.5	84.2	1.14	30.1	4.5
All-Tex 7A21	1843	44.7	84.2	1.16	29.6	4.5
PHY 375 WRF	1838	43.8	84.3	1.13	29.0	4.1
BX 1262B2F	1808	44.0	83.7	1.17	30.4	4.3
CG 3787 B2RF	1805	45.7	85.2	1.17	27.7	4.2
DP 1133 B2RF	1774	45.6	85.7	1.17	32.4	4.5
AM 1550 B2RF	1710	43.5	83.7	1.12	27.6	4.6
ST 4288B2F	1709	40.0	84.2	1.17	30.2	4.6
All-Tex ATX3039 B2RF	1708	43.8	84.1	1.14	27.9	3.9
BRS286	1687	41.2	84.6	1.13	30.6	4.5
ST 4145LLB2	1666	43.5	84.5	1.15	30.3	4.3
DP 0924 B2RF	1628	43.2	84.4	1.15	28.2	4.6
BRS293	1625	40.0	84.3	1.15	32.2	4.4
SSG CT310 HQ	1569	39.2	84.5	1.16	34.5	4.0
GA2004143	1567	43.9	85.0	1.20	33.1	4.2
DP 0949B2RF	1566	43.9	84.1	1.19	31.5	5.0
SSG HQ 210 CT	1552	41.6	84.3	1.18	29.7	4.3
BX 1252LLB2	1549	41.0	85.2	1.16	32.3	4.0
All-Tex LA122	1541	43.9	84.4	1.15	27.6	4.3
FM1740B2RF	1518	43.5	83.9	1.17	29.9	4.4
GA2008057	1493	41.4	85.0	1.20	31.4	4.2
BCSX 1150B2RF	1430	40.1	85.6	1.23	33.4	4.2
All-Tex ATX81144	1426	41.3	85.2	1.25	31.1	3.9
PHY 367 WRF	1425	43.7	83.9	1.16	29.8	4.2
GA2006106	1401	42.2	84.9	1.18	33.8	4.5
SSG CT Linwood	1348	43.4	83.6	1.09	33.2	4.8
Average	1684	43.2	84.5	1.16	30.6	4.4
LSD 0.10	305	0.8	1.1	0.05	3.0	0.4
CV %	15.4	1.6	0.8	2.60	5.7	5.5

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: April 27, 2011.

Harvested: September 13, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 78 lb N, 54 lb P₂O₅, and 108 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Rainfall (in):	1.48	0	1.94	4.06	1.26	4.53

Trials conducted by Larry Thompson.

Yield Summary for Dryland Earlier Maturity Cotton Varieties, 2011

Variety	Lint Yield ^a					Lint %	Unif. Index %	Length in	Strength g/tex	Mic. units
	Athens	Midville	Plains lb/acre	Tifton	4-Loc. Average					
PHY 499 WRF	1725 ³	1238 ¹	604 ¹⁰	2067 ²	1409 ¹	46.2	83.9	1.10	31.1	4.4
AM1511 B2RF	1778 ¹	1123 ²	648 ³	1946 ⁵	1374 ²	45.5	83.8	1.09	30.3	4.5
DP 0912 B2RF	1544 ⁷	1116 ³	687 ¹	1964 ⁴	1328 ³	43.5	83.8	1.12	29.9	4.7
DP 1028 B2RF	1395 ¹³	1039 ⁵	672 ²	2130 ¹	1309 ⁴	47.2	84.1	1.13	27.9	4.5
AM 1550 B2RF	1678 ⁴	997 ⁹	606 ⁹	1710 ¹²	1248 ⁵	43.6	83.7	1.10	26.8	4.4
BX 1262B2F	1773 ²	1002 ⁸	381 ³⁰	1808 ⁹	1241 ⁶	42.6	83.3	1.13	31.2	4.1
All-Tex 7A21	1607 ⁶	989 ¹¹	475 ²⁷	1843 ⁷	1228 ⁷	44.1	83.7	1.13	30.1	4.5
Dyna-Gro 2570B2RF	1450 ¹¹	961 ¹⁴	553 ¹⁷	1916 ⁶	1220 ⁸	42.6	83.7	1.11	29.6	4.4
ST 4288B2F	1663 ⁵	982 ¹³	500 ²⁵	1709 ¹³	1214 ⁹	40.8	83.5	1.13	28.0	4.3
CG 3787 B2RF	1463 ¹⁰	994 ¹⁰	581 ¹²	1805 ¹⁰	1211 ¹⁰	46.3	84.2	1.13	28.1	4.5
DP 0920 B2RF	1321 ²²	840 ²⁴	607 ⁸	1986 ³	1189 ¹¹	43.8	83.8	1.13	26.9	4.3
DP 0924 B2RF	1505 ⁹	1011 ⁷	528 ²¹	1628 ¹⁷	1168 ¹²	42.4	83.7	1.10	28.4	4.5
ST 4145LLB2	1371 ¹⁶	1063 ⁴	518 ²³	1666 ¹⁶	1155 ¹³	43.6	83.0	1.11	29.8	4.2
PHY 375 WRF	1341 ²¹	949 ^{16T}	476 ²⁶	1838 ⁸	1151 ¹⁴	44.1	83.4	1.09	28.3	4.1
BRS293	1512 ⁸	986 ¹²	459 ²⁸	1625 ¹⁸	1146 ¹⁵	41.4	83.1	1.10	32.9	4.7
All-Tex ATX3039 B2RF	1433 ¹²	939 ¹⁷	431 ²⁹	1708 ¹⁴	1128 ¹⁶	44.6	82.5	1.11	26.7	4.1
BRS286	1306 ²⁵	859 ²³	614 ⁷	1687 ¹⁵	1116 ¹⁷	41.0	82.9	1.09	30.3	4.3
DP 0949B2RF	1357 ¹⁷	949 ^{16T}	566 ¹⁵	1566 ²¹	1109 ^{18T}	44.5	83.3	1.12	30.0	4.7
BX 1252LLB2	1376 ¹⁵	886 ²⁰	627 ⁵	1549 ²³	1109 ^{18T}	42.1	83.6	1.12	30.9	4.3
All-Tex LA122	1355 ¹⁹	916 ¹⁸	578 ¹³	1541 ²⁴	1098 ¹⁹	44.4	83.4	1.11	27.8	4.3
DP 1133 B2RF	1240 ²⁸	793 ²⁶	542 ¹⁸	1774 ¹¹	1087 ^{20T}	45.3	84.2	1.13	30.7	4.6
GA2004143	1352 ²⁰	874 ²¹	555 ¹⁶	1567 ²⁰	1087 ^{20T}	45.2	83.8	1.15	31.9	4.4
PHY 367 WRF	1318 ²³	958 ¹⁵	645 ⁴	1425 ²⁹	1086 ²¹	43.9	83.6	1.12	29.7	4.2
All-Tex ATX81144	1356 ¹⁸	1036 ⁶	511 ²⁴	1426 ²⁸	1082 ²²	41.3	84.7	1.21	32.3	3.7
SSG HQ 210 CT	1313 ²⁴	862 ²²	588 ¹¹	1552 ²²	1079 ²³	41.4	82.6	1.10	30.7	4.6
BCSX 1150B2RF	1383 ¹⁴	888 ¹⁹	534 ¹⁹	1430 ²⁷	1059 ²⁴	40.4	84.1	1.17	31.8	4.1
FM1740B2RF	1241 ²⁷	816 ²⁵	567 ¹⁴	1518 ²⁵	1035 ²⁵	43.2	83.2	1.12	28.5	4.2
GA2006106	1253 ²⁶	754 ^{28T}	618 ⁶	1401 ³⁰	1006 ²⁶	41.9	83.6	1.15	31.9	4.3
SSG CT310 HQ	1166 ²⁹	754 ^{28T}	519 ²²	1569 ¹⁹	1002 ²⁷	39.9	83.4	1.12	33.4	4.5
SSG CT Linwood	872 ³¹	761 ²⁷	531 ²⁰	1348 ³¹	878 ²⁸	43.4	82.9	1.08	31.9	4.9
GA2008057	1004 ³⁰	621 ²⁹	317 ³¹	1493 ²⁶	859 ²⁹	41.0	84.3	1.16	32.8	4.3
Average	1402	934	550	1684	1142	43.3	83.6	1.12	30.0	4.4
LSD 0.10	254	128	N.S. ^b	305	144	1.2	0.7	0.02	1.3	0.2
CV %	15.4	11.7	31.1	15.4	17.2	1.9	0.9	2.19	4.7	5.1

^a Superscripts indicate ranking at that location.

^b The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Two-Year Summary for Dryland Earlier Maturity Cotton Varieties at Four Locations^a, 2010-2011

Variety	Lint Yield lb/acre	Lint %	Uniformity Index %	Length inches	Strength g/tex	Micronaire units
PHY 499 WRF	1303	46.2	83.5	1.09	31.2	4.6
DP 1028 B2RF	1194	47.2	83.3	1.11	28.1	4.8
DP 0912 B2RF	1187	43.7	82.9	1.08	29.2	4.9
PHY 375 WRF	1072	44.5	82.9	1.08	27.9	4.3
Dyna-Gro 2570B2RF	1067	43.0	83.1	1.09	29.4	4.6
AM 1550 B2RF	1055	43.4	83.0	1.08	26.7	4.5
DP 0920 B2RF	1052	44.2	82.9	1.11	26.7	4.5
All-Tex 7A21	1042	44.3	83.3	1.13	29.9	4.7
All-Tex LA122	1033	44.3	83.2	1.11	27.9	4.5
DP 0924 B2RF	1026	43.1	82.7	1.07	28.1	4.7
ST 4288B2F	1023	40.8	82.4	1.10	27.4	4.5
PHY 367 WRF	1020	43.7	82.7	1.11	29.2	4.3
FM1740B2RF	977	43.3	82.6	1.09	28.5	4.5
GA2006106	966	41.9	82.8	1.15	31.6	4.5
SSG CT Linwood	864	43.1	82.3	1.07	31.4	5.0
Average	1059	43.8	82.9	1.10	28.9	4.6
LSD 0.10	73	0.4	0.4	0.02	0.7	0.1
CV %	16.8	2.0	0.9	2.54	4.5	5.3

^a Athens, Midville, Plains, and Tifton.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Athens, Georgia:

Dryland Later Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
PHY 499 WRF	1515	49.4	84.4	1.09	31.2	4.9
DP 1048 B2RF	1434	48.9	83.3	1.07	28.6	4.9
BX 1262B2F	1411	46.3	83.4	1.10	30.1	4.7
DP 1137 B2RF	1407	48.2	83.8	1.11	28.2	5.0
DP 1050 B2RF	1365	48.6	83.5	1.11	27.5	4.6
ST 5458B2RF	1320	45.0	82.6	1.07	30.2	5.0
DP 1034 B2RF	1284	47.9	83.3	1.10	27.4	4.7
BX 1254LLB2	1276	46.3	82.9	1.13	29.6	4.7
AM1511 B2RF	1226	47.9	83.5	1.07	29.6	4.5
MON 10R051 B2RF	1178	49.5	83.0	1.07	27.4	4.9
DP 1252 B2RF	1169	49.7	84.0	1.12	27.4	5.1
ST 4145LLB2	1158	45.9	82.1	1.08	31.2	4.1
ST 5288B2F	1123	44.6	83.2	1.08	27.6	4.8
BX 1252LLB2	1115	45.7	83.9	1.13	33.2	4.5
PHY 440 W	1114	45.4	84.1	1.09	30.9	4.4
GA2007095	1075	43.3	83.0	1.10	29.8	4.3
PHY 565 WRF	1075	44.6	83.8	1.07	30.0	4.3
DP 1133 B2RF	1062	47.7	83.3	1.08	30.3	5.2
GA2004230	1046	44.0	84.2	1.15	30.1	4.4
PHY 375 WRF	987	47.1	83.0	1.07	28.1	4.5
GA2008083	973	49.0	82.7	1.05	32.7	4.6
BX 1261B2F	892	42.8	83.2	1.09	29.9	4.5
Average	1191	46.7	83.3	1.09	29.6	4.6
LSD 0.10	195	0.8	1.0	0.04	1.7	0.4
CV %	13.9	1.4	0.7	2.35	3.4	4.7

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 5, 2011.

Harvested: September 19, 2011.

Soil Type: Cecil coarse sandy loam.

Fertilization: 180 lb N, 80 lb P₂O₅, and 80 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Rainfall (in):	2.50	1.40	3.69	1.90	1.00	1.63

Trials conducted by Larry Thompson.

Midville, Georgia:
Dryland Later Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
PHY 499 WRF	1168	47.0	82.9	1.07	29.8	4.7
AM1511 B2RF	1150	45.3	83.0	1.05	29.3	4.7
BX 1262B2F	1146	46.6	81.5	1.08	28.7	4.8
ST 5458B2RF	1134	44.3	81.3	1.06	26.4	4.8
BX 1261B2F	1082	43.5	82.6	1.10	29.9	4.1
DP 1137 B2RF	1056	47.0	84.3	1.10	28.6	4.8
ST 5288B2F	1042	44.9	82.8	1.06	27.0	4.4
DP 1048 B2RF	1040	46.0	82.4	1.06	27.0	4.5
PHY 565 WRF	1039	44.2	83.8	1.10	30.0	4.5
DP 1034 B2RF	1032	45.1	82.4	1.08	26.6	4.8
DP 1050 B2RF	1025	46.7	83.3	1.11	26.7	4.9
BX 1254LLB2	1014	44.0	82.3	1.12	30.1	4.7
BX 1252LLB2	1007	43.6	83.5	1.12	30.0	4.7
MON 10R051 B2RF	1007	46.5	82.7	1.10	27.8	4.9
GA2004230	1003	44.0	82.5	1.15	29.2	4.9
ST 4145LLB2	995	44.0	82.4	1.09	29.0	4.5
PHY 375 WRF	976	45.8	81.7	1.08	27.6	4.4
GA2007095	952	43.7	82.6	1.13	30.2	4.8
PHY 440 W	862	43.4	83.4	1.10	32.3	4.3
DP 1252 B2RF	840	46.0	82.7	1.07	27.8	4.8
DP 1133 B2RF	840	46.6	82.9	1.06	31.2	5.1
GA2008083	725	44.1	82.4	1.07	30.5	4.8
Average	1006	45.1	82.7	1.09	28.9	4.7
LSD 0.10	163	2.1	1.2	0.04	2.2	0.3
CV %	13.7	3.9	0.8	2.37	4.4	3.3

* A random quality sample was taken on the picker during cotton harvest.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 3, 2011.

Harvested: October 3, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 85 lb N, 60 lb P₂O₅, and 70 lb K₂O/acre.

Management: Temik applied 5 lb/acre and Telone II applied 3 gal/acre.

	April	May	June	July	Aug.	Sept.
Irrigation (in):	1.41	2.32	1.76	2.22	1.50	2.51

Trials conducted by Larry Thompson.

Plains, Georgia:
Dryland Later Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
AM1511 B2RF	762	42.6	82.3	1.11	31.0	4.2
DP 1050 B2RF	680	41.9	82.4	1.11	27.7	3.8
PHY 440 W	650	37.8	82.9	1.11	30.4	3.5
DP 1252 B2RF	646	42.7	83.2	1.11	30.0	3.9
DP 1048 B2RF	632	41.7	83.0	1.12	28.0	3.8
MON 10R051 B2RF	628	41.9	82.4	1.11	28.1	3.8
PHY 375 WRF	627	40.5	82.5	1.08	26.7	3.6
GA2007095	625	37.5	82.9	1.15	30.9	3.8
DP 1137 B2RF	625	41.4	82.6	1.09	27.8	4.2
PHY 499 WRF	594	44.8	82.8	1.10	32.0	4.1
DP 1034 B2RF	591	40.4	83.2	1.13	28.4	3.7
ST 4145LLB2	576	38.1	82.8	1.13	29.4	3.7
GA2004230	571	37.9	82.5	1.19	30.0	4.1
BX 1262B2F	569	40.5	82.4	1.13	31.7	3.9
ST 5288B2F	543	38.3	82.0	1.10	27.7	3.9
ST 5458B2RF	535	38.5	81.8	1.11	30.0	4.3
DP 1133 B2RF	508	41.5	83.6	1.13	31.4	4.0
GA2008083	500	42.6	81.9	1.14	32.3	4.2
PHY 565 WRF	480	35.9	82.4	1.08	29.1	3.5
BX 1254LLB2	449	39.0	81.0	1.12	29.7	3.6
BX 1261B2F	388	37.0	82.4	1.12	29.8	3.7
BX 1252LLB2	386	39.1	82.4	1.14	31.1	3.8
Average	571	40.1	82.5	1.12	29.7	3.8
LSD 0.10	156	1.4	N.S. ¹	N.S.	2.7	0.3
CV %	23.1	3.1	1.5	3.11	5.2	5.0

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: May 5, 2011.

Harvested: October 7, 2011.

Soil Type: Greenville sandy clay loam.

Fertilization: 100 lb N, 88 lb P₂O₅, and 24 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Rainfall (in):	2.01	0.03	1.12	7.00	1.50	3.38

Trials conducted by Larry Thompson.

Tifton, Georgia:
Dryland Later Maturity Cotton Variety Performance, 2011

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length* inches	Strength* g/tex	Micronaire* units
			Index* %			
PHY 499 WRF	2254	45.2	84.6	1.18	32.6	4.3
DP 1252 B2F	2022	47.5	84.9	1.16	27.3	4.7
ST 5288B2F	1957	43.7	84.3	1.15	26.2	4.4
DP 1137 B2RF	1945	44.3	85.2	1.15	29.3	4.6
DP 1133 B2RF	1940	46.3	85.2	1.15	31.1	4.6
BX 1262B2F	1906	43.6	84.6	1.17	30.7	4.7
AM1511 B2RF	1906	46.0	85.0	1.15	29.7	5.0
DP 1050 B2RF	1905	45.3	84.2	1.16	26.5	4.5
BX 1254LLB2	1895	44.6	84.2	1.11	32.0	4.9
MON 10R051 B2RF	1850	46.7	85.1	1.16	28.2	4.4
DP 1034 B2RF	1844	46.6	84.9	1.15	27.0	4.5
DP 1048 B2RF	1814	44.9	85.5	1.18	28.1	4.4
BX 1261B2F	1690	40.5	83.8	1.16	28.7	4.2
PHY 375 WRF	1676	43.2	84.4	1.16	29.1	3.9
PHY 565 WRF	1658	41.3	83.5	1.19	31.2	4.1
GA2004230	1657	42.5	84.3	1.23	30.5	4.1
ST 4145LLB2	1650	43.5	84.5	1.15	31.3	4.5
ST 5458B2RF	1626	42.3	83.5	1.17	32.4	5.0
BX 1252LLB2	1593	42.1	84.0	1.19	31.1	4.2
GA2007095	1558	42.2	84.7	1.19	28.1	4.2
PHY 440 W	1536	41.6	84.6	1.12	30.6	4.4
GA2008083	1442	46.6	83.1	1.09	31.1	4.3
Average	1787	44.1	84.4	1.16	29.7	4.4
LSD 0.10	242	0.7	N.S. ¹	0.04	1.9	0.3
CV %	11.5	1.3	1.0	2.17	3.8	3.7

* A random quality sample was taken on the picker during cotton harvest.

1. The F-test indicated no statistical differences at the alpha = .10 probability level; therefore a LSD value was not calculated.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

Planted: April 27, 2011.

Harvested: September 13, 2011.

Soil Type: Tifton loamy sand.

Fertilization: 78 lb N, 54 lb P₂O₅, and 108 lb K₂O/acre.

Management: Temik applied 5 lb/acre.

	April	May	June	July	Aug.	Sept.
Rainfall (in):	1.48	0	1.94	4.06	1.26	4.53

Trials conducted by Larry Thompson.

Yield Summary for Dryland Later Maturity Cotton Varieties, 2011

Variety	Lint Yield ^a					Lint %	Unif. Index %	Length in	Strength g/tex	Mic. units
	Athens	Midville	Plains lb/acre	Tifton	4-Loc. Average					
PHY 499 WRF	1515 ¹	1168 ¹	594 ⁹	2254 ¹	1383 ¹	46.6	83.6	1.11	31.4	4.5
AM1511 B2RF	1226 ⁹	1150 ²	762 ¹	1906 ^{6T}	1261 ²	45.4	83.4	1.09	29.9	4.6
DP 1137 B2RF	1407 ⁴	1056 ⁶	625 ^{8T}	1945 ⁴	1258 ^{3T}	45.2	83.9	1.11	28.5	4.6
BX 1262B2F	1411 ³	1146 ³	569 ¹³	1906 ^{6T}	1258 ^{3T}	44.3	83.0	1.12	30.3	4.5
DP 1050 B2RF	1365 ⁵	1025 ¹¹	680 ²	1905 ⁷	1244 ⁴	45.6	83.3	1.12	27.1	4.4
DP 1048 B2RF	1434 ²	1040 ⁸	632 ⁵	1814 ¹¹	1230 ⁵	45.4	83.5	1.11	27.9	4.4
DP 1034 B2RF	1284 ⁷	1032 ¹⁰	591 ¹⁰	1844 ¹⁰	1188 ⁶	45.0	83.4	1.12	27.3	4.4
DP 1252 B2RF	1169 ¹¹	840 ^{19T}	646 ⁴	2022 ²	1169 ⁷	46.5	83.7	1.11	28.1	4.6
ST 5288B2F	1123 ¹³	1042 ⁷	543 ¹⁴	1957 ³	1166 ^{8T}	42.9	83.1	1.10	27.1	4.4
MON 10R051 B2RF	1178 ¹⁰	1007 ^{13T}	628 ⁶	1850 ⁹	1166 ^{8T}	46.1	83.3	1.11	27.9	4.5
BX 1254LLB2	1276 ⁸	1014 ¹²	449 ¹⁹	1895 ⁸	1159 ⁹	43.5	82.6	1.12	30.3	4.5
ST 5458B2RF	1320 ⁶	1134 ⁴	535 ¹⁵	1626 ¹⁷	1154 ¹⁰	42.5	82.3	1.10	29.7	4.7
ST 4145LLB2	1158 ¹²	995 ¹⁵	576 ¹¹	1650 ¹⁹	1094 ¹¹	42.9	82.9	1.11	30.2	4.2
DP 1133 B2RF	1062 ¹⁷	840 ^{19T}	508 ¹⁶	1940 ⁵	1087 ¹²	45.5	83.7	1.10	31.0	4.7
GA2004230	1046 ¹⁸	1003 ¹⁴	571 ¹²	1657 ¹⁵	1069 ¹³	42.1	83.4	1.18	29.9	4.3
PHY 375 WRF	987 ¹⁹	976 ¹⁶	627 ⁷	1676 ¹³	1067 ¹⁴	44.2	82.9	1.10	27.8	4.1
PHY 565 WRF	1075 ^{16T}	1039 ⁹	480 ¹⁸	1658 ¹⁴	1063 ¹⁵	41.5	83.3	1.11	30.1	4.1
GA2007095	1075 ^{16T}	952 ¹⁷	625 ^{8T}	1558 ¹⁹	1052 ¹⁶	41.7	83.3	1.14	29.7	4.3
PHY 440 W	1114 ¹⁵	862 ¹⁸	650 ³	1536 ²⁰	1041 ¹⁷	42.1	83.7	1.10	31.0	4.1
BX 1252LLB2	1115 ¹⁴	1007 ^{13T}	386 ²¹	1593 ¹⁸	1025 ¹⁸	42.6	83.5	1.14	31.3	4.3
BX 1261B2F	892 ²¹	1082 ⁵	388 ²⁰	1690 ¹²	1013 ¹⁹	41.0	83.0	1.12	29.6	4.1
GA2008083	973 ²⁰	725 ²⁰	500 ¹⁷	1442 ²¹	910 ²⁰	45.6	82.5	1.09	31.7	4.5
Average	1191	1006	571	1787	1139	44.0	83.2	1.11	29.4	4.4
LSD 0.10	195	163	156	242	134	1.1	0.6	0.02	1.2	0.2
CV %	13.9	13.7	23.1	11.5	14.3	2.6	1.0	2.53	4.2	4.2

^a Superscripts indicate ranking at that location.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

**Two-Year Summary for Dryland Later Maturity
Cotton Varieties at Four Locations^a, 2010-2011**

Variety	Lint Yield lb/acre	Lint %	Uniformity	Length inches	Strength g/tex	Micronaire units
			Index %			
PHY 499 WRF	1267	46.5	83.4	1.09	31.3	4.6
DP 1050 B2RF	1144	46.2	83.3	1.11	27.4	4.6
DP 1137 B2RF	1143	45.6	83.4	1.10	27.9	4.7
DP 1048 B2RF	1115	45.8	83.1	1.10	27.7	4.6
DP 1034 B2RF	1081	45.4	83.1	1.11	27.6	4.6
DP 1252 B2RF	1063	46.9	83.3	1.10	28.4	4.8
ST 5458B2RF	1040	42.8	82.0	1.09	29.3	4.8
ST 5288B2F	1030	43.0	82.6	1.09	27.2	4.6
DP 1133 B2RF	1019	45.8	83.1	1.09	31.1	4.8
PHY 375 WRF	991	44.4	82.5	1.08	28.1	4.3
PHY 565 WRF	963	41.8	82.9	1.10	30.5	4.4
PHY 440 W	938	43.3	83.0	1.07	30.3	4.3
Average	1066	44.8	83.0	1.09	28.9	4.6
LSD 0.10	62	0.4	0.5	0.02	0.6	0.1
CV %	14.0	2.4	0.9	2.65	3.7	5.0

^a Athens, Midville, Plains, and Tifton.

Bolding indicates entries not significantly different from highest yielding entry based on Fisher's protected LSD (P = 0.10).

TOBACCO

Tifton, Georgia:

Official Flue-Cured Tobacco Variety Test -

Yield, Value, Price Index, Grade Index, and Agronomic and Chemical Characteristics of Released Varieties, 2011

Variety	Yield	Value	Price Index ¹	Grade Index ²	Leaves/ Plant	Plant Ht.	Days to Flower	Total Alkaloids	Reducing Sugars	Ratio RS/TA
	lb/A	\$/A	\$/CWT		number	in		%	%	
NC 196	3639	4405	121	65	20	38.9	80	2.77	16.4	5.94
GF 318	3632	4808	134	53	19	36.7	75	2.30	18.6	8.11
CC 37	3581	3998	112	61	19	39.6	79	2.15	17.4	8.06
NC 92	3564	3155	88	50	21	40.1	80	3.19	14.5	4.54
NC 291	3554	4619	128	69	20	36.2	74	2.89	16.1	5.58
SP 168	3498	4426	125	67	19	36.7	76	1.96	16.6	8.47
PVH 1452	3474	4201	121	65	20	37.1	75	2.69	16.5	6.14
SP 227	3398	4102	118	64	20	37.1	79	2.08	14.8	7.12
CC 65	3369	3216	94	54	20	42.1	79	2.62	15.4	5.89
PVH 2277	3367	5182	154	79	19	39.1	71	2.17	17.7	8.15
CC 700	3355	4409	131	69	20	37.3	73	3.15	14.9	4.72
NC 71	3345	4387	131	70	18	35.2	78	2.43	15.1	6.22
NC 297	3321	4094	124	67	19	35.3	79	3.15	16.2	5.16
GL 395	3301	4314	129	70	20	37.9	79	2.39	15.1	6.31
NC 299	3289	4000	121	67	19	37.0	82	2.85	15.8	5.54
PVH 1596	3277	4030	124	68	19	35.7	76	3.18	13.7	4.31
GL 338	3274	3875	118	54	18	35.8	72	2.70	16.8	6.22
CC 27	3262	3892	120	64	18	36.1	72	2.82	13.0	4.62
K 399	3231	4019	122	66	20	34.8	74	2.41	18.0	7.45
NC 72	3228	3562	111	62	19	36.3	79	2.67	13.9	5.22
NC 471	3211	4267	134	72	19	39.8	78	2.71	15.2	5.62
SP 225	3155	4155	132	71	18	36.5	77	2.34	15.8	6.77
K 346	3114	3166	101	55	18	34.7	78	2.61	14.5	5.56
K 326	3075	5372	173	89	18	35.1	75	2.31	13.5	5.87
SP 236	3007	3464	111	61	19	35.5	79	2.21	16.7	7.56
GL 368	2951	3010	98	70	19	38.3	79	2.51	15.9	6.33
CC 67	2939	3386	111	61	18	37.1	75	1.93	18.7	9.71
NC 95	2725	4417	162	82	18	38.3	73	2.67	14.1	5.27
NC2326	2154	2249	105	53	17	34.2	64	2.56	13.8	5.37
LSD @ 0.05	616.3	1500.4	32.6	14.5						

Conducted on an Ocilla loamy sand soil fertilized with 1100 lbs/a of 6-6-18 and 163 lbs/a 15.5-0-0 with plants spaced 20-22 inches apart in 44-inch rows.

1. Price Index based on two year average (2010-2011) prices for U.S. government grades.
2. Numerical values ranging from 1-99 for flue-cured tobacco based on equivalent government grades - higher the number, higher the grade.

Researched by Stevan S. LaHue and C.E. Troxell supported by grants from the Georgia Tobacco Commission.

Tifton, Georgia:
Three and Two -Year Averages of Official Flue-Cured Tobacco
Variety Test - Comparison of Released Varieties
for Certain Characteristics, 2008, 2010 and 2011

Variety	Yield	Value	Price Index ¹	Grade Index ²	Leaves/ Plant	Plant Ht.	Days to Flower	Total Alkaloids	Reducing Sugars	Ratio RS/TA
	lb/A	\$/A	\$/CWT		number	in		%	%	
3 Year Average 2008, 2010 and 2011										
NC 196	3436	4878	143	73	21	40	79	2.33	16.3	7.12
Speight 227	3408	4808	140	71	20	38	77	2.55	15.0	6.03
NC 291	3367	4594	136	71	20	36	76	2.95	14.6	5.03
CC 37	3361	4288	129	66	18	40	78	2.55	15.2	6.17
NC 71	3360	4825	144	72	20	35	75	2.54	15.3	6.06
NC 92	3345	4247	127	65	20	40	77	2.85	15.3	5.45
Speight 168	3298	4605	141	71	19	37	75	2.40	15.5	6.65
NC 72	3262	4445	137	70	21	38	75	2.87	13.8	4.87
CC 27	3252	4295	133	68	21	39	73	2.52	14.1	5.70
NC 297	3211	4563	142	71	20	38	76	2.70	16.0	6.02
K 326	3206	5053	157	78	21	38	79	2.62	15.0	4.95
CC 700	3164	4522	142	72	20	38	75	2.82	15.5	5.53
Speight 236	3102	4308	137	70	20	38	76	2.73	16.0	6.03
Speight 225	3051	4477	147	74	19	38	75	2.51	15.1	6.02
K 346	3051	4173	136	69	21	36	76	2.55	14.0	5.48
NC 299	2994	4259	143	72	20	38	80	2.57	16.2	6.38
NC 95	2741	3857	141	72	19	39	75	3.04	14.5	4.91
NC 2326	2407	2786	116	59	18	37	66	3.19	12.5	4.25
2 Year Average 2010-2011										
GF 318	3398	4826	144	64	20	40	76	2.67	17.8	6.87
CC 37	3319	4195	128	67	16	40	79	2.27	16.2	7.15
PVH 1452	3286	4501	138	71	19	38	74	2.64	15.4	5.85
NC 196	3213	4473	142	72	20	40	80	2.48	16.4	6.72
SP 227	3142	4206	134	69	19	36	79	2.58	14.3	5.80
NC 92	3140	3482	114	61	20	40	77	2.77	14.5	5.37
SP 168	3139	4314	139	71	18	36	76	2.27	15.8	7.13
NC 291	3131	4254	135	71	19	35	76	2.75	15.5	5.65
NC 471	3101	4507	146	75	20	41	78	2.49	15.0	6.04
CC 27	3081	3945	129	67	20	38	74	2.48	13.8	5.69
NC 72	3068	3981	132	69	19	38	77	2.63	14.1	5.36
GL 338	3044	4067	135	64	18	37	70	2.70	15.9	5.90
CC 700	3031	4192	139	71	19	38	75	2.94	15.7	5.40
PVH 1596	2973	4119	141	73	19	37	75	2.69	15.3	6.02
NC 71	2950	4123	141	73	19	35	77	2.40	15.2	6.34
SP 236	2912	3772	128	67	19	37	75	2.70	16.4	6.30
K326	2892	4618	158	80	19	36	78	2.63	14.2	5.48
K 399	2892	4017	140	72	19	35	77	2.35	17.8	7.54
NC 297	2877	3853	136	70	19	36	78	2.74	16.1	6.02
PVH 2277	2851	4477	158	79	19	37	75	2.36	17.8	7.59

**Three and Two -Year Averages of Official Flue-Cured Tobacco
Variety Test - Comparison of Released Varieties
for Certain Characteristics, 2008, 2010 and 2011 (Continued)**

Variety	Yield	Value	Price Index ¹	Grade Index ²	Leaves/ Plant	Plant Ht.	Days to Flower	Total Alkaloids	Reducing Sugars	Ratio RS/TA
	lb/A	\$/A	\$/CWT		number	in		%	%	
2 Year Average 2010-2011 - continued										
NC 299	2838	3841	138	71	19	37	82	2.50	15.6	6.32
K346	2808	3529	128	67	18	36	75	2.52	14.4	5.72
SP 225	2808	4011	144	74	18	37	78	2.44	15.4	6.33
GL 368	2791	3590	128	74	18	38	76	2.75	15.8	5.81
CC 67	2703	3480	128	67	18	36	76	2.12	17.3	8.29
NC 95	2452	3707	150	75	19	39	77	2.68	15.1	5.62
NC2326	2278	2931	127	64	18	36	66	2.51	12.8	5.08

spaced 20-22 inches apart in 44-inch rows.

1. Price Index based on two year average (2010-2011) prices for U.S. government grades.

2. Numerical values ranging from 1-99 for flue-cured tobacco based on equivalent government grades - higher the number, higher the grade.

Researched by Stevan S. LaHue and C.E. Troxell supported by grants from the Georgia Tobacco Commission.


Tifton, Georgia:
Regional Farm Flue-Cured Tobacco Variety Test -
Comparison of Varieties for Certain Characteristics, 2011

Variety	Yield	Value	Price Index ¹	Grade Index ²	Leaves/ Plant number	Plant Ht. in	Days to Flower	Total Alkaloids %	Reducing Sugars %	Ratio RS/TA
	lb/A	\$/A	\$/CWT							
NCEX 34	3772	5798	155	81	19	39.3	72	2.00	14.0	7.01
CU 136	3722	5823	157	81	19	39.5	70	2.18	14.5	6.64
GLEX 328	3720	6079	163	84	19	38.9	76	1.99	15.2	7.61
CC 1063	3685	5969	162	84	19	38.6	70	2.80	12.7	4.53
ULT 123	3590	6000	167	86	19	38.9	73	1.76	15.6	8.86
ULT 143	3580	5889	164	85	18	38.0	71	2.24	14.5	6.45
NC EX 24	3502	5338	153	80	18	39.9	74	1.91	13.7	7.18
NC 95	3390	4876	144	74	19	41.1	73	2.67	14.1	5.27
NCTG 156	3345	5500	164	85	18	34.3	69	2.07	13.8	6.64
PXH 1	3336	5035	150	78	20	39.1	77	1.93	13.6	7.08
XP 254	3275	5229	165	86	19	40.0	75	1.84	15.8	8.61
RJR 901	3176	4933	155	82	19	38.3	73	2.39	11.3	4.73
K 326	3161	5236	164	85	19	36.5	75	2.31	13.5	5.87
GF 157	2920	4901	167	86	18	37.3	70	2.17	11.1	5.13
GLEX 362	2766	4405	159	84	18	35.9	73	2.40	12.8	5.34
NC 2326	2070	2674	129	67	17	34.6	64	2.56	13.8	5.37
LSD -0.05	571.1	1105.8	16.9	7.53						

Conducted on an Ocilla loamy sand soil fertilized with 1100 lbs/a of 6-6-18 and 163 lbs/a 15.5-0-0 with plants spaced 20-22 inches apart in 44-inch rows.

1. Price Index based on two-year average (2010-2011) prices for U.S. government grades.
2. Numerical values ranging from 1-99 for flue-cured tobacco based on equivalent government grades - higher the number, higher the grade.

Researched by Stevan S. LaHue and C.E. Troxell supported by grants from the Georgia Tobacco Commission.



★ Main Experiment Station ★ Branch Station ● Extension Center

University of Georgia

Agricultural Experiment Stations
Athens, Georgia 30602
Robert Shulstad, Associate Dean

Publication

Penalty for Private Use \$300

ADDRESS CORRECTION REQUESTED

The University of Georgia and Ft. Valley State University, the U.S. Department of Agriculture and counties of the state cooperating, Cooperative Extension, and the University of Georgia College of Agricultural and Environmental Sciences offer educational programs, assistance and materials to all people without regard to race, color national origin, age, gender or disability.

**An Equal Opportunity Employer/Affirmative Action Organization
Committed to a Diverse Work Force**

“CERTIFIED SEED DOESN’T COST ... IT PAYS”

HERE’S WHY:

- Known performance of varieties adapted to your area.
- A pedigree record that begins with the release of breeder seed and continues until it reaches the consumer as certified (blue tag) seed.
- Field inspected for trueness to variety and inseparable from other crop and weed seed.
- Certified seed can only be conditioned in an approved facility.
- Certified seed must meet High Quality standards as to germination and purity.
- Free of noxious weeds.

The planting of CERTIFIED SEED eliminates many of the risks associated with crop production. For sources of Certified seed, contact your local county Extension agent or the Georgia Crop Improvement Association, Inc. (706-542-2351)

